

PREVALENCE OF VITAMIN D DEFICIENCY IN DIABETIC PATIENTS

Taha Mahwi ^a and Rawand M. Raouf ^b

Submitted: 6/3/2018; Accepted: 5/8/2018; Published 15/10/2018



ABSTRACT

Background

There are increasing incidence of diabetes around the worldwide, and there are a lot of studies have been considering vitamin D's role in the incidence, and improving glucose status in diabetes.

Objectives

This study is done to evaluate the prevalence of vitamin D deficiency in diabetic patients in Sulaimani Diabetes and Endocrine Center.

Materials and Methods

This is a case-control study done on 100 diabetic patients compared to 100 persons in the general population in Sulaimani/Iraq. Their Vitamin D was estimated from June 1st, 2017 to December 1st, 2017.

Results

Results: The mean age of the diabetic group was (49.68 yr. \pm 8.64 SD), and the mean age of the non-diabetic group was (50.49 yr. \pm 6.78 SD). The mean vitamin D's level in the diabetic group was (16.91 ng/dl \pm 5.5 SD) and in a non-diabetic group was (16.21 ng/dl \pm 8.59) (P-values= 0.493). Among diabetic group vitamin D level is significantly higher in males with an average of 6.2 ng/dl in comparison to females, (P-Value = 0.024), among diabetic patients, Vitamin D level was deficient in (66%), insufficient in (19%) and was sufficient just in (15%), while in non-diabetic control group vitamin D level was deficient in (77%), insufficient in (20%) and sufficient just in (3%). The P-value for these differences is (0.0052).

Conclusion

The results of this study show the majority of our participants have suboptimal vitamin D level. Vitamin D level was lower in female patients in comparison to male patients.

Keywords: *Diabetes Mellitus (DM), Vitamin D deficiency, Serum Vitamin D.*

^a Department of Medicine, College of Medicine, University of Sulaimani.

Correspondence: rawandraouf@gmail.com

^b Kurdistan Board of Medical Specialties Candidate

INTRODUCTION

Diabetes mellitus (DM) has become one of a large public health problem, affecting more than 300 million individuals in the world, with significant complications worldwide ⁽¹⁾ with the increasing number of patients with DM all over the world. It is expected that this disorder will be a reason for significant morbidity and mortality ⁽²⁾.

Due to lifestyle changes and amount of food intake after the discovery of oil in the 1950s in Arabian Gulf countries the numbers of diabetes and metabolic syndrome are increasing at an alarming rate ^(3,4).

Vitamin D inadequacy constitutes a largely unrecognized epidemic in many populations worldwide, postmenopausal ^(5,6), persistent non-specific musculoskeletal pain ⁽⁷⁾, primary hyperparathyroidism in female patients ⁽⁸⁾, hip fracture in elderly females ⁽⁹⁾, osteoarthritis in postmenopausal females ⁽¹⁰⁾, secondary hyperparathyroidism ⁽¹¹⁾, minimal trauma causing fracture ⁽¹²⁾ and osteoporosis ⁽¹³⁾. A study done in Karballa/Iraq showed that vitamin D insufficiency and deficiency was highly prevalent in postmenopausal women which was more than (85%), women of child-bearing age was more than (65%), young men aged between (25-49) years was more than (60%) and men aged between (50-70) years was about (82%) ⁽¹⁴⁾.

Despite musculoskeletal defects, low vitamin D is related to some non-skeletal disorders such as malignancy, autoimmune diseases, hypertension, endothelial dysfunction, dyslipidemia, infections, and cardiovascular disease ⁽¹⁵⁾.

More recently, many studies were done to prove the effect of vitamin D deficiency and calcium status in the development of type 2 DM ^(16,17), most of them may have a role ⁽¹⁸⁻²¹⁾, but not all of them ^(22,33).

In recent years, studies have done showed that using vitamin D can help to decrease the incidence of diabetes mellitus and adjustment of insulin and glucose ⁽²³⁻²⁷⁾. Surveys showed that using supplements of vitamin D with calcium will significantly decrease fasting blood glucose level ⁽²⁸⁾. So, using vitamin D supplements alone can prevent type-1 diabetes incidence ⁽²⁹⁾.

There is no survey showing prevalence of vitamin D deficiency in diabetic patients in our region, this study is done to evaluate prevalence and vitamin D status in diabetic patients.

PATIENTS AND METHODS

This is a case-control study done in Sulaimani city. Hundred patients with type 2 diabetes mellitus, randomly selected in Sulaimani Diabetes and Endocrine Center to evaluate vitamin D level from June 1st, 2017 to December 1st, 2017, after they had agreed to participate in this study with inclusion criteria. The diabetic patients with renal insufficiency and who received vitamin D supplement are excluded and set as a case group. The control group selected from the general population matched with gender and sex to case group.

Serum vitamin D sent for all the patients and a case group fasting blood sugar, HbA1c, renal function and lipid profile was evaluated for diabetic group, due to seasonal change effect on vitamin D, all vitamin D was collected in one season. Vitamin D was measured with Cobas electrochemiluminescence immunoassay (ECLIA) on Cobas e411. Vitamin D was defined:

- Deficient when serum 25-hydroxyvitamin D is less than 20 ng/ml.
- Insufficient when serum 25-hydroxyvitamin D is between 20-30 ng/ml.
- Sufficient when serum 25-hydroxyvitamin D is concentration higher than 30 ng/dl (30).

Statistical analysis

Vitamin D level was dealt with as an independent variable and HbA1c, fasting blood sugar, lipid profile, Body Mass Index was dealt with as dependent variables in the statistical analysis. Data management and analysis were performed utilizing Statistical Package for Social Sciences (SPSS) software, version 24.

This study was done on human participants. All data were kept in a secure and confidential manner. All participants' identification and associated data were separated. All data were analyzed as total population in a manner that individual privacy was maintained. All records, results and progress, both electronic and composed will be kept with the researchers in case of review.

A written informed consent was taken from the participants clarifying the aim of the study, the importance of the respondents' views, the researcher's name.

RESULTS

Prevalence of Vitamin D Deficiency in Diabetic Patients

In this study, one hundred (100) patients with Diabetes Mellitus had been screened for vitamin D deficiency, and they were compared to (100) non-diabetic individuals randomly selected from the general population set as the control group.

Both groups are well matched in terms of age (patients with type 2 diabetes mean Age: 49.68 yr. \pm 8.64 SD vs. Non-diabetics mean Age: 50.49 yrs. \pm 6.78 SD), P-Value (0.461), that is no significant difference in age. The same for gender (diabetics; 33% male, 67% female), control (39% male, 61% female), P-Value: (0.465).

The mean concentration of vitamin D in diabetic group was (16.91 ng/dl \pm 5.5 SD) and in non-diabetic patients was (16.21 ng/dl \pm 8.59 SD) (P-values= 0.493).

Among the diabetic group, vitamin D level is significantly higher in males by an average of 6.2 ng/dl in comparison to females, (p-Value: 0.024). While this difference is not present in non-diabetics (male 16.98 \pm 5.02 Vs. female 16.87 \pm 5.82). (P-Value = 0.12), Table 1.

Among diabetic patients, Vitamin D level was deficient in (66%) and insufficient in (19%) and was sufficient in (15%), while in the non-diabetic control group, vitamin D level was deficient in (77%), insufficient in (20%), and sufficient in (3%). The P-Value for these differences is (0.0052). Figure 1.

Among diabetic group, rate of deficiency was higher in females, 75% of females were deficient (n=50), (15%) (n=10) were insufficient, and (10%) (n=7) were sufficient. While male patients had (48%) rate of deficiency (n=16), (27%) (n=9) were insufficient and (25%) (n=8) were sufficient. The p-value is (0.056), Table 2.

A weak negative correlation between serum vitamin D level and HbA1C% was found by a factor of (- 0.182) and the P-value for this correlation is 0.06.

Also, the negligible correlation between serum Vitamin D level and BMI was found by a factor of (- 0.018) and the P-value for this correlation is 0.8.

Table 1. The Case and control group's mean age with vitamin D levels

	Diabetic group	Non-diabetic group	P value
Mean Vitamin D	16.91 \pm 5.5 SD	16.21 \pm 8.59 SD	0.493
Mean Age	49.68 yrs. \pm 8.64 SD)	50.49 yrs. \pm 6.78 SD	0.461
Male(n)	33	39	0.465
Female (n)	67	61	
Male Vitamin D	22.40 \pm 10.73	16.98 \pm 5.02	p-value 0.12
Female Vitamin D	16.21 \pm 8.59	16.87 \pm 5.82	




Figure 1. Prevalence of vitamin D deficiency in the diabetic and non-diabetic group.

Table 2. Vitamin D deficiency in diabetic according to Gender.

	Females	Males	P-Value
Deficient	50 (75%)	16 (48%)	0.056
Insufficient	10 (15%)	9 (27%)	
Sufficient	7 (10%)	8 (25%)	
Total (n)	67(100%)	33 (100)	

DISCUSSION

The result of this study shows a very high prevalence of vitamin D deficiency in both diabetic and non-diabetic group but the mean vitamin D level in both groups was near to each other (P-value was 0.493) which is considered not a significant difference between diabetic and non-diabetic. In other studies done in Iran and Saudi Arabia, there was a significant difference seen in the diabetic group, vitamin D was lower in the diabetic group compared to the non-diabetic group^(31, 32). The difference in the results may be due to that our study has the smaller sample size and the age of our diabetic and non-diabetic group was between (37-64) years, which means participant's age was middle age and elderly patients in this study. In two other studies, one of them was done in Iraq to know the variation of vitamin D in respect of age; sex and nationality both

prove that vitamin D level affected by age, sex and gender^(33, 34).

The vitamin D's level was higher among male patients than female patients with significant differences; P-value of this was (0.024). This could be explained by that females have less sun exposure compared to males, and wear more clothes that cover most parts of the body and go out little.

There is a negative relationship between vitamin D and HbA1c which means those diabetic patients have a low level of vitamin D, have poor control of their glycemic control (P-Value = 0.06). This need to be proved by the correction of vitamin D and excluding other causes of poor glycemic control. In the studies done to prove that, most but not all, showed that correction of vitamin D affects glycemic control⁽³⁵⁾.

Conclusions

The results of this study show that the great majority of our participants have suboptimal vitamin D level. In female patients, vitamin D level was lower than in male patients.

REFERENCES

1. Sherwin, R.; Jastreboff, A.M. Year in diabetes 2012: the diabetes tsunami. *J. Clin. Endocrinol. Metab.* 2012, 97,4293–4301
2. Zella JB, DeLuca HF. Vitamin D and autoimmune diabetes. *J Cell Biochem* 2003; 88: 216-22.
3. Motlagh B, O'Donnell M, Yusuf S. Prevalence of cardiovascular risk factors in the Middle East: a systematic review. *Eur J Cardiovasc Prev Rehabil.* 2009;16(3):268–80.
4. Al Zenki S, Al Omirah H, Al Hooti S, Al Hamad N, Jackson RT, Rao A, et al. High prevalence of metabolic syndrome among Kuwaiti adults—a wake-up call for public health intervention. *Int J Environ Res Public Health.* 2012;9(5):1984–96.
5. Gaugris S, Heaney RP, Boonen S, Kurth H, Bentkover JD, Sen SS. Vitamin D inadequacy among post-menopausal women: a systematic review. *QJM.* 2005;98:667-676.
6. Isaia G, Giorgino R, Rini GB, Bevilacqua M, Maugeri D, Adami S. Prevalence of hypovitaminosis D in elderly women in Italy: clinical consequences and risk factors. *Osteoporos Int.* 2003;14:577-582.
7. Plotnikoff GA, Quigley JM. Prevalence of severe hypovitaminosis D in patients with persistent, nonspecific musculoskeletal pain. *Mayo Clin Proc.* 2003;78:1463-1470
8. Carnevale V, Manfredi G, Romagnoli E, et al. Vitamin D status in female patients with primary hyperparathyroidism: does it play a role in skeletal damage? *Clin Endocrinol (Oxf).* 2004;60:81-86
9. Harwood RH, Sahota O, Gaynor K, Masud T, Hosking DJ. A randomized, controlled comparison of different calcium and vitamin D supplementation regimens in elderly women after hip fracture: the Nottingham Neck of Femur (NONOF) Study. *Age Ageing.* 2004;33:45-51
10. Glowacki J, Hurwitz S, Thornhill TS, Kelly M, LeBoff MS. Osteoporosis and vitamin-D deficiency among postmenopausal women with osteoarthritis undergoing total hip arthroplasty. *J Bone Joint Surg* 11. Gomez-Alonso C, Naves-Diaz ML, Fernandez-Martin JL, Diaz-Lopez JB, Fernandez-Coto MT, Cannata-Andia JB. Vitamin D status and secondary hyperparathyroidism: the importance of 25-hydroxyvitamin D cut-off levels. *Kidney Int Suppl.* 2003;85: S44-S48.
12. Blau EM, Brenneman SK, Bruning AL, Chen Y. Prevalence of vitamin D insufficiency in an osteoporosis population in Southern California [abstract]. *J Bone Miner Res.* 2004;19(suppl 1): S342. Abstract SU582.
13. Simonelli C, Weiss TW, Morancey J, Swanson L, Chen YT. Prevalence of vitamin D inadequacy in a minimal trauma fracture population. *Curr Med Res Opin.* 2005;21:1069-1074.
14. Al-Hilali, K. A.; Prevalence of Hypovitaminosis D in Adult Iraqi People Including Postmenopausal Women; *Scientific Research Journal (SCIRJ)*, Volume IV, Issue IX, September 2016
15. Binkley N, Ramamurthy R, Krueger D. Low vitamin D status: definition, prevalence, consequences, and correction. *Endocrinol Metab Clin North Am.* 2010;39:287-301
16. Bischoff-Ferrari HA, Giovannucci E, Willett WC, Dietrich T, Dawson-Hughes B. Estimation of optimal serum concentrations of 25-hydroxyvitamin D for multiple health outcomes. *Am J Clin Nutr* 2006; 84:18–28.
17. Holick MF. High prevalence of vitamin D inadequacy and implications for health. *Mayo Clin Proc* 2006; 81:353–373.
18. Cigolini M, Iagulli MP, Miconi V, Galiotto M, Lombardi S, Targher G. Serum 25-hydroxyvitamin D3 concentrations and prevalence of cardiovascular disease among type 2 diabetic patients. *Diabetes Care* 2006; 29:722–724.
19. Hypponen E, Power C. Vitamin D status and glucose homeostasis in the 1958 British birth cohort: the role of obesity. *Diabetes Care* 2006; 29:2244–2246.
20. Aksoy H, Akcay F, Kurtul N, Baykal O, Avci B. Serum 1,25 dihydroxy vitamin D (1,25(OH)2D3), 25 hydroxyvitamin D (25(OH)D) and parathormone levels in diabetic retinopathy. *Clin Biochem* 2000; 33:47–51.
21. Isaia G, Giorgino R, Adami S. High prevalence of hypovitaminosis D in female type 2 diabetic population. *Diabetes Care* 2001; 24:1496.

22. Ishida H, Seino Y, Matsukura S, Ikeda M, Yawata M, Yamashita G, et al. Diabetic osteopenia and circulating levels of vitamin D metabolites in type 2 (noninsulin-dependent) diabetes. *Metabolism* 1985; 34:797-801
23. Lambert PW, Service FJ, Arnaud SB. Calcium homeostasis in diabetes mellitus. *J Clin Endocrinol Metab* 1979; 49:462-466.
24. Zipitis CS, Akobeng AK. Vitamin D supplementation in early childhood and risk of type 1 diabetes: a systematic review and meta-analysis. *Arch Dis Child* 2008; 93: 512-7.
25. Boursolon PM, Billaudel B, Faure-Dussert A. Influence of vitamin D3 deficiency and 1, 25 dihydroxy vitamin D3 on de novo insulin biosynthesis in the islets of the rat endocrine pancreas. *J Endocrinol* 1999; 160: 87-95.
26. Baynes KC, Boucher BJ, Feskens EJ, Kromhout D. Vitamin D, glucose tolerance and insulinemia in elderly men. *Diabetologia* 1997; 40: 344-7.
27. Pittas AG, Harris SS, Stark PC, Dawson-Hughes B. The effects of calcium and vitamin D supplementation on blood glucose and markers of inflammation in nondiabetic adults. *Diabetes Care* 2007; 30: 980-6.
28. Stene LC, Ulriksen J, Magnus P, Joner G. Use of cod liver oil during pregnancy associated with lower risk of Type I diabetes in the offspring *Diabetologia* 2000; 43:1093-8.
29. Tangpricha V, Pearce EN, Chen TC, Holick MF. Vitamin D insufficiency among free-living healthy young adults. *Am J Med* 2002; 112: 659-62.
30. Daga RA, Laway BA, Shah ZA, et al. High prevalence of vitamin D deficiency among newly diagnosed youth-onset diabetes mellitus in north India. *Arq Bras Endocrinol Metabol* 2012; 56: 423-8.
31. Bayani MA, Banasaz B, Saeedi F. Status of Vitamin-D in diabetic patients and compared it with healthy individuals. *Caspian J Intern Med* 2014; 5(1): 40-42.
32. Al-Zaharani M. Prevalence of Vitamin D Deficiency in type 2 DM patients. *Majmaah J. Health Sciences* 2013;1(1): 24-28
33. Al-Horani, Hanan et al. Nationality, Gender, Age, and Body Mass Index Influences on Vitamin D Concentration among Elderly Patients and Young Iraqi and Jordanian in Jordan. *Biochemistry Research International* 2016 (2016): 8920503
34. Sinem BOZKURT, Berat Meryem ALKAN, Fatma YILDIZ, et al. Age, Sex, and Seasonal Variations in the Serum Vitamin D3 Levels in a Local Turkish Population. *Turkish League Against Rheumatism*, 2014; 29(1): 014-019
35. Madar AA, Knutsen KV, Stene LC, et al. Effect of vitamin D3 supplementation on glycated hemoglobin (HbA1c), fructosamine, serum lipids, and body mass index: a randomized, double-blinded, placebo-controlled trial among healthy immigrants living in Norway. *BMJ Open Diabetes Research and Care* 2014;2:e000026.