



Serological Detection of SAT-2 Serotype Foot and Mouth Disease Virus in Buffalo, Iraq

Thaer W. Enad ✉

Khalefa A. Mansour

Department of Internal and Preventive Veterinary Medicine, College of Veterinary Medicine, University of Al-Qadisiyah, Iraq

<https://orcid.org/0000-0001-9135-7256>

Submitted: July 24, 2025
Revised: December 9, 2025
Accepted: December 26, 2025

Correspondence

Thaer W. Enad
Vet.post22.10@qu.edu.iq

Abstract This study aims to determine the SAT-2 serotype causing FMD outbreak in buffalo in Wassit and Dhi-Qar governorates, Iraq. This was done by clinical and serological methods. ELISA technology, which measures antibody levels in buffalo serum samples. 92 blood samples were collected from infected buffalo for antibody screening against FMDV serotype SAT-2 in infected animals which develop antibodies against the non-structural proteins (NSP) of the virus. Affected animals showed various clinical signs, such as fever, decreased appetite, mouth and foot vesicles, and lameness. According to the results of serological tests, the infection rate reached 85/92 positive samples (92.39%). The results showed, Age and sexspecific investigations of FMDV seropositivity did not reveal any statistically significant differences. The age groups (6-18 months and 18-36 months) had lowest infection rate (91.42%) and (92%) respectively, whereas the age group 36 months, had highest rate of infection at (93.75%). Also, the results revealed, the seroprevalence rates infection for males were (90.6%), which had lowest than females (93.33%). This study, confirms a widespread outbreak of FMD caused by the SAT2 serotype in buffalo populations across the Wassit and Dhi-Qar governorates of Iraq. The study's findings may offer helpful information about how FMD protection measures are established and carried out in Iraq.

Keywords: Non-structural proteins, ELISA, SAT-2, Buffalo, FMD

©Authors, 2025, College of Veterinary Medicine, University of Al-Qadisiyah. This is an open access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction One of the most important infectious diseases that can harm all animals with cloven hooves is FMD(1). This is because epidemics typically have significant economic impacts, including both direct livestock losses and indirect losses due to the high cost of containment and trade restrictions that often occur in the aftermath of an FMD outbreak, which is caused by the aphthovirus of the Picornaviridae family (2,3). The disease impairs the international transport traffic of live animals and their by-products, presenting a serious global danger.

Seven serotypes of FMDV are known to be present:” A, O, C, Asia1, SAT-1, SAT-2, and SAT-3’(4,5)Each genotype exhibits distinct subtypes that differ

antigenically as a result of a high mutation rate (4,6). FMD is widespread in numerous countries on the African continent, Asia, and Latin America (7,8). In Iraq, FMDV serotypes Asia 1, A, and O are constantly present and lead to occasional epidemics (9,10). In 2023, significant outbreaks caused by FMDV serotype SAT-2 occurred in the provinces of Iraq (11). All serotypes of FMD virus induce the same clinical symptoms(12,13), but immunization to one serotype does not protect others due to their broad antigenic variability (14). Through either direct or indirect contact with sick animals,

contaminated materials, or through the air, the disease can spread rapidly (15). Buffalo that are affected typically exhibit fever and discontinue ruminating. Saliva was shaped like a ruby. blisters on the lips, mouth, tongue, nose, and between the toes. The amount of milk produced has decreased (16,17). The infected animals can be discriminated from vaccinated by non-structural proteins (NSP) which can be detected only in infected animals (18). ELISA is an easy and efficient technique compared to virus isolation and RT-PCR. for high-throughput testing that does not require extensive technical expertise or specialised equipment (19,20). Early in 2023, animals all over Iraq were affected by a large FMD outbreak (21). The primary aim of the research was to determine the FMD virus SAT-2 Serotype present in Iraqi buffalo.

material and Methods

Ethical approvals

This study has been accepted by the Scientific Committee of the College of Veterinary Medicine at Al-Qadisiyah University in Iraq.

Collection of samples

Samples for this investigation were taken from 92 water buffalo that were suspected to have FMD. After a clinical examination, the buffalo displayed certain clinical signs. Using a sterile syringe, from each sick animal, 10 milliliters of blood was taken from the jugular vein, using a gel tube, and the sample was left for 30 minutes to clot. at 3000 rpm for 10 minutes collect the serum, transfer it to Eppendorf tubes (22), and store the samples at -20°C (23). The samples were properly labeled and transported to the laboratory under refrigeration for further processing.

Sandwich Enzyme-Linked Immunosorbent Assay (ELISA)

A total of 92 animals who exhibited symptoms of FMD underwent a ten-milliliter blood sample collection procedure, it was tested using an ELISA reaction to see if the non-structural (NSP) antibody was specific to FMDV serotype SAT-2.

Result

Clinical Results

In our study, we observed clinical signs suggestive of FMD, such as fever, dullness, and loss of appetite. Also, the result of clinical study revealed the infected animals suffering from, drooling saliva and hanging in long ropy strings up to the ground (**Figure 1-A**). Erosions, and ulcers on the nostrils and muzzle (**Figure 1-B**). The mouths of infected buffalos suggestive lesions of FMD. The gums, dental pad, hard palate, tongue, and

interior of the oral cavity contained ruptured vesicles, erosions, or ulcers (**Figure 2-A**), and skin lesions that were develop in the coronary region and interdigital cavity of the hoof (**Figure 2-B**).

Figure 1: Shows clinical signs of FMD, A: saliva



hanging in long ropy strings, B: nostril lesion



Figure 2: Shows clinical signs of FMD, A: The dental pad, hard palate, and interior of the oral cavity contained ruptured vesicles, erosions, or ulcers B: lesion around the coronary band.

Serological Results

The prevalence of FMD seropositivity in buffalo of both sexes and ages. Of the 92 buffalo shown 85 had positive FMDV SAT-2 antibodies, Figures 2 and 3, indicating that 92.39% of them were infected. According to the results of ELISA, the age groups of buffalo (6-18 months and 18-36 months) had lowest rate of infection affected with FMD respectively, with 32/35 (91.42 %), 23/25 (92%). as shown as in Figure (2) FMD, whereas the highest incidence was found among third group of age at 30/32 (93.75) but does not have statistical significance differences at ($p > 0.05$).

The effect of sex on FMDV seropositivity was examined. Figure (3) shows the

seropositivity rates in males and females. The results showed that males had a seropositivity rate of 29/32 (90.6%) while females had a higher seropositivity rate of 56/60 (93.33%). However, the disparity in seropositivity rates among males and females did not reach significance in statistics ($p > 0.05$).

structural proteins such as 3AB, 3ABC, 2C, and other targets. Antibody occurred on days 5–10 post-infection in all animals with no history of prior exposure to FMDV antigens(26,27).

Infected animals develop antibodies against nonstructural proteins (NSP) of the virus. In contrast, because vaccines are inactivated and purified, recipient animals only receive antibodies against structural proteins from vaccines(28)(29,30)

The results of serological tests showed the presence of antibodies to FMDV virus in 92.39% of the samples, indicating a high prevalence of the disease among the study population.

The high seropositivity in our results from this study is consistent with the high seropositivity reported in Al-Qadisiyah and Basrah (31,32). while the rates of infection in Al-Najaf and Diyala were (34%, 25.33%)(33,34). The high

seropositivity rate observed in our study may be because all samples were collected at the optimal time from animals infected with a novel strain not previously registered in Iraq Moreover, the inclusion of animals lacking vaccination against this particular strain may have further contributed to the observed high seropositivity rate.

The infection rate of FMD according to age the results of this study, no statistically significant findings were discovered regarding the age of identified cases that agreed with other studies(35,36), that showed statistically no significant difference between age in seroprevalence of FMD.

The lack of significant differences in the high incidence of infection between different age groups might be due to the attribute of rearing by not separating young animals from their mothers as well as the communal grazing practices(34).

Although higher incidence of females than males, no statistically significant findings were discovered regarding the gender of identified cases. The results of this study agreed with many other epidemiological studies on FMD The results of this study agreed with many other epidemiological studies on FMD which found a slight increase in female infection (37,38) This may be attributed to physiological factors like lactation, pregnancy, and estrus(32,39)

The gender prevalence of FMDV appears to contradict previous reports from others that showed increased




Figure (2). FMD serology results in Buffalo across various Age (months)groups




Figure (3) FMD Serology results in Buffalo across various Sex groups

The clinical examination of the buffalo in this study revealed the presence of typical signs of FMD, including fever, reduced appetite, ulcers in the mouth and foot, and impaired mobility. These signs are consistent with previous studies and provide further evidence of the presence of FMDV in the studied population(24,25). This serological investigation aimed to determine the serotype of the FMD virus responsible for the recent epidemic in livestock in Iraq.

Differentiation between vaccinated and infected animals is important for prevention planning. ELISA is used to detect infected animals by detecting non-



seropositivity in male animals compared with female animals.

Identifying serotypes is very important, as it allows the implementation of appropriate prevention and control strategies(42).

The results of our study are important for developing and implementing plans to manage FMD in Iraq because the high prevalence of SAT-2 serotype in buffalo emphasizes the need for vaccination campaigns targeted against serotype spread throughout the country.

In this study, the use of ELISA as a diagnostic tool to detect FMDV antibodies in buffalo serum samples showed its effectiveness and reliability(43).

In summary, this investigation has shown that the present FMD outbreak in Iraq is caused by serotype SAT2 and this serotype's high prevalence highlights the need for targeted immunization.

Conclusion

The results of our studies have proven the spread of the Sat -2 serotype for the first time in Iraq. The high prevalence of FMDV antibodies in the buffalo studied indicates the need for Iraq. The high prevalence of FMDV antibodies in the buffalo studied indicates the need for targeted vaccination campaigns, specifically the SAT-2 serotype. The use of ELISA as a diagnostic tool proves to be effective and reliable in identifying FMDV antibodies in serum samples from buffalo.

Conflict of interest

There is no conflict of interest in this study as stated by the authors.

Acknowledgment

Not applicable.

Funding source

This research had no specific fund; however, it was self-funded by the authors.

References

1. Iriarte M V., Gonz ales JL, de Freitas Costa E, Gil AD, de Jong MCM. Main factors associated with foot-and-mouth disease virus infection during the 2001 FMD epidemic in Uruguay. *Front Vet Sci.* 2023;10.
2. Kabir A, Kamboh AA, Ahmed S, Marri NU. Foot Mouth Disease Virus Affects The Immune Response Of The Host : An Overview. 2024;11(1):18–26.
3. Mkama M, Philemon W, Donald K, David P, Francois M. Challenges of controlling foot-and-mouth disease in pastoral settings in Africa. 2024;2024(1):1–12.

4. Anjume H, Hossain KA, Hossain A, Hossain MA. Heliyon Complete genome characterization of foot-and-mouth disease virus My-466 belonging to the novel lineage O / ME-SA / SA-2018. *Heliyon* [Internet]. 2024;(December 2023):e26716. Available from: <https://doi.org/10.1016/j.heliyon.2024.e26716>
5. Byomi¹ A, Zidan¹ S, Sakr¹ E, Eissa¹ N, Elsobky¹ Y. Journal of Current Veterinary Research Epidemiological Patterns of Foot and Mouth Disease in Egypt and Other African Countries. 2023; Available from: <http://www.jcivr.journals.ekb.eg>
6. Shin SH, Hwang SY, Kim HM, Shin SH, Ko MK, Lee MJ, et al. Evaluation of a Vaccine Candidate Designed for Broad-Spectrum Protection against Type A Foot-and-Mouth Disease in Asia. *Vaccines.* 2024;12(1):1–12.
7. Nishi T, Fukai K, Morioka K. Molecular Basis of the Pathogenicity of the Foot-and-Mouth Disease Virus Isolated in Japan. *Japan Agric Res Q.* 2024;58(1):25–30.
8. Junior AAF, Laguardia-Nascimento M, Ferreira APS, da Silva RR, J nior AVR, Camargos MF. Comparison of seven different RT-qPCR for diagnostic of Foot-and-Mouth disease virus. *Cienc Anim Bras.* 2024;25.
9. Aslam M, Alkheraije KA. The prevalence of foot-and-mouth disease in Asia. *Front Vet Sci.* 2023 Jun 30;10.
10. Sheikh MB, Rashid PA, Raheem Z, Marouf AS, Amin KM. Molecular characterization and phylogenetic analysis of foot and mouth disease virus isolates in Sulaimani province, Iraq. *Vet Res Forum.* 2021;12(2):247–51.
11. FAO World Reference Laboratory for Foot-and-Mouth Disease (WRLFMD) Genotyping Report [Internet]. Available from: [www:http://www.wrlfmd.org/](http://www.wrlfmd.org/)
12. El Damaty HM, Fawzi EM, Neamat-Allah ANF, Elsohaby I, Abdallah A, Farag GK, et al. Characterization of foot and mouth disease virus serotype sat-2 in swamp water buffaloes (*Bubalus bubalis*) under the egyptian smallholder production system. *Animals.* 2021;11(6).
13. Paton DJ, Di Nardo A, Knowles NJ, Wadsworth J, Pituco EM, Cosivi O, et al. The history of foot-and-mouth disease virus serotype C: The first known extinct serotype? *Virus Evol.* 2021;7(1):1–12.
14. Zewdie G, Akalu M, Tolossa W, Belay H, Deresse G, Zekarias M, et al. A review of foot-and-mouth disease in Ethiopia: epidemiological aspects, economic implications, and control strategies. *Virology* [Internet]. 2023;20(1):299. Available from: <https://doi.org/10.1186/s12985-023-02263-0>
15. Brown E, Nelson N, Gubbins S, Colenutt C. Airborne Transmission of Foot-and-Mouth Disease



- Virus: A Review of Past and Present Perspectives. *Viruses*. 2022;14(5):1–14.
16. Mohammed MM. The study of prevalence foot and mouth disease in Al-Muthana province. *AL-Qadisiyah J Vet Med Sci*. 2013;12(2).
17. Humadi AA. MICROBIAL AND VIRULENT VIEW TO CAUSATIVE AGENT OF FOOT AND MOUTH DISEASE IN UNVACCINATED DAIRY COWS IN DIYALA CAJMNS. 2024;(c):8–11.
18. Hosamani M, Basagoudanavar SH, Tamil Selvan RP, Das V, Ngangom P, Sreenivasa BP, et al. A multi-species indirect ELISA for detection of non-structural protein 3ABC specific antibodies to foot-and-mouth disease virus. *Arch Virol*. 2015;160(4):937–44.
19. Cao Y, Li K, Xing X, Zhu G, Fu Y, Bao H, et al. Development and Validation of a Competitive ELISA Based on Bovine Monoclonal Antibodies for the Detection of Neutralizing Antibodies against Foot-and-Mouth Disease Virus Serotype A. *J Clin Microbiol*. 2022;60(4).
20. Tewari A, Jain B, Bhatia AK. Multiplexed DIVA tests for rapid detection of FMDV infection/circulation in endemic countries. *Appl Microbiol Biotechnol*. 2020;104(2):545–54.
21. Qualitative FAO, Assessment R. Risk of foot-and-mouth disease SAT2 introduction and spread in countries in the Near East and West Eurasia. Risk of foot-and-mouth disease SAT2 introduction and spread in countries in the Near East and West Eurasia. 2023.
22. Alkefari OA, Al-Gharban HA, Ahmed TH. Microscopic, serological and molecular detection of *Babesia bigemina* in buffaloes (*Bubalus bubalis*) in Wasit province, Iraq. *Al-Qadisiyah J Vet Med Sci*. 2017;16(1):123–30.
23. Al-abodi HRJ. Serological and molecular detection of *Toxoplasma gondii* in *Columba livia* hunting pigeons of AL-Qadisiyah province. *Al-Qadisiyah J Vet Med Sci*. 2017;16(1):136–41.
24. Ismail I, Indarjulianto S, Yusuf S, Purba FY. Clinical Examination of Foot and Mouth Disease of Dairy Cows in Sukamurni, Cilawu, Garut, West Java, Indonesia. *IOP Conf Ser Earth Environ Sci*. 2023;1174(1).
25. Ulaiwi HK, Al-Tamemy HMJ, Mansoor NH. Histopathological alteration of lung, small intestine and lymph nodes in calves suffering from typical clinical case of Foot and Mouth Disease. *J Vet Med Sci*. 2012;38(2).
26. Zia MA, Dobson SJ, Rowlands DJ, Stonehouse NJ, Shah MS, Habib M. Development of an ELISA to distinguish between foot-and-mouth disease virus infected and vaccinated animals utilising the viral non-structural protein 3ABC. *J Med Microbiol*. 2022;71(4):1–12.
27. Health C. The value of diagnostic assays in FMD control. 2017;(October).
28. Subramaniam Saravanan RR. Diagnosis of Foot and Mouth Disease | ResearchGate. . 2012;(February).
29. Clavijo A, Sanchez-Vazquez MJ, Buzanovsky LP, Martini M, Pompei JC, Cosivi O. Current Status and Future Prospects to Achieve Foot-and-Mouth Disease Eradication in South America. *Transbound Emerg Dis*. 2017;64(1):31–6.
30. Colling A, Morrissy C, Barr J, Meehan G, Wright L, Goff W, et al. Development and validation of a 3ABC antibody ELISA in Australia for foot and mouth disease. *Aust Vet J*. 2014;92(6):192–9.
31. A Ajeel Y, AN Alordhan M, H Kshash Q. Clinical and Diagnostic Study of Foot and Mouth Disease by using ELISA test in Cattle of AL-Diwaniya. *AL-Qadisiyah J Vet Med Sci*. 2012;11(2):91–8.
32. Al-Rodhan AM. Detection of Cattle Foot and Mouth disease Virus by RT-PCR and ELISA. *Kufa Journal For Veterinary Medical Sciences*.
33. Abd Hatem A, Ahmed Abdul Wahid Al Anbagi N, Al-Alo KZK, Sabah Bustani G. Detection of Antibody against Non-Structural Proteins of Foot-and-Mouth Disease Virus in Cattle in Najaf, Iraq. *Arch Razi Inst*. 2022;77(3):1185–9.
34. Sadun Al-Ajeeli K, Khazal Al-Azawy A, Khudhair AL-Anbagi G, Mohammed Salih Abdul-Rasoul L. Sero-Prevalence of Foot-and-Mouth Disease in Cattle by 3ABC NSP ELISA *Address for Correspondence [Internet]. Vol. 9, *Indian Journal of Natural Sciences* www.tnsroindia.org.in ©IJONS. 2018. Available from: www.tnsroindia.org.in©IJONS
35. Dubie T, Negash W. Seroprevalence of bovine foot and mouth disease (FMD) and its associated risk factors in selected districts of Afar region, Ethiopia. *Vet Med Sci*. 2021;7(5):1678–87.
36. Salim SAS, Talb AOQ, Yousif AS, Daher HS. Prevalence and risk factors of foot and mouth disease virus in Nineveh province, Iraq. *Adv Anim Vet Sci*. 2020 Jan 1;8(1):1–10.
37. Belina D, Girma B, Mengistu S. Sero-Prevalence of Bovine Foot and Mouth Disease in Selected Districts of Eastern Showa Zone , Oromia Regional State , Ethiopia SeroPrevalenceofBovineFootandMouthDiseaseinSelectedDistrictsofEasternShowaZoneOromiaRegionalStateEthiopia. 2016;(July).
38. Atuman YJ, Kudi CA, Abdu PA, Okubanjo OO, Abubakar A, Wungak Y, et al. Seroprevalence of Foot and Mouth Disease Virus Infection in Some Wildlife and Cattle in Bauchi State, Nigeria. *Vet Med Int*. 2020;2020.
39. Aghaalikhani, Hossein & Ahmadi E. Research Article Research Article. *Arch Anesthesiol Crit Care* [Internet]. 2018;4(4):527–34. Available from:



<http://www.globalbuddhism.org/jgb/index.php/jgb/article/view/88/100>

40. Chowdhury MSR, Ahsan MI, Khan MJ, Rahman MM, Hossain MM, Harun-Al-Rashid A, et al. Data on prevalence, distribution and risk factors for Foot and Mouth Disease in grazing cattle in haor areas of Bangladesh. *Data Br* [Internet]. 2020;28:104843. Available from:

<https://doi.org/10.1016/j.dib.2019.104843>

41Hasan M, Mia MM. Prevalence, Distribution and Epidemiological Risk Factors of Foot-and-Mouth

Disease Among the Cattle Population of Bangladesh: a Meta-Analysis. *Bulg J Vet Med.* 2023;26(4):542–55.

42. Das S, Pal S, Rautaray SS, Mohapatra JK, Subramaniam S, Rout M, et al. Estimation of foot-and-mouth disease virus sero-prevalence rates using novel computational approach for the susceptible bovine population in India during the period 2008–2021. *Sci Rep* [Internet]. 2023;13(1):1–17. Available from: <https://doi.org/10.1038/s41598-023-48459-w>

43. Wong CL, Yong CY, Ong HK, Ho KL, Tan WS. Advances in the Diagnosis of Foot-and-Mouth Disease. *Front Vet Sci.* 2020;7(August):1–24.