



Effect of different extraction methods on apigenin yield, antioxidant activity, and anticoagulant potential of watercress (*Eruca vesicaria* L.) leaves

Nabaa Qais Jameel

Department of Biology, University Education for Women, University of Anbar, Anbar, Iraq.

* Corresponding author: E-mail: nabaa.qais@uoanbar.edu.iq

ABSTRACT

Watercress (*Eruca vesicaria*) is a rich source of secondary metabolites with disease-preventive properties, including the phenol apigenin. This work aimed to examine the impact of various extraction methods on apigenin yield from watercress (*Eruca vesicaria* L.) leaves and its antioxidant capacity/blood coagulant effect. Organisations compared modern with traditional extraction methods. Active compounds identification and overall qualitative chemical analysis proved that the extract is rich in flavonoids, phenols, and terpenes. The selected solvent and the method of extraction had a remarkable impact on both yield and purity of apigenin. Free radical scavenging ability of saliva and phenolic content were also found high and these values were correlated, suggesting that its phenols may significantly contribute for the high oxidative activity. The results of this study further showed that apigenin is involved in modulating the blood coagulation system via prolonging prothrombin time (PT) and activating activated partial thromboplastin time (aPTT). This indicates that Ap inhibits intrinsic as well extrinsic coagulation paths. These results suggest that the leaves of *E. vesicaria* represent a rich source of apigenin, which may be used as potential phytochemicals in nutraceutical and cardiovascular disorders related to oxidative stress and thrombosis-induced diseases prevention.

KEYWORDS: Apigenin's estimation, different extraction techniques, oxidative activity, clotting factors, PT, PTT.

Received: 04/02/2026; Accepted: 26/03/2026; Available online: 31/03/2026

©2026. This is an open access article under the CC by licenses <http://creativecommons.org/licenses/by/4.0>

التأثير المقارن لتقنيات الاستخلاص على إنتاجية الأبيجينين من أوراق نبات الجرجير البري (*Eruca vesicaria* L.): القدرة المضادة للأكسدة وتعديل التخثر


نبأ قيس جميل

قسم علوم الحياة، كلية التربية للبنات، جامعة الأنبار، الانبار، العراق

المخلص

يُعد نبات الجرجير المائي (*Eruca vesicaria*) مصدراً غنياً بالمركبات الأيضية الثانوية ذات الخصائص الوقائية من الأمراض، بما في ذلك الفينول أبيجينين. هدفت هذه الدراسة إلى فحص تأثير طرق الاستخلاص المختلفة على إنتاجية الأبيجينين من أوراق الجرجير المائي (*Eruca vesicaria* L.) وقدرته المضادة للأكسدة وتأثيره على تخثر الدم. قارنت الدراسة بين طرق الاستخلاص الحديثة والتقليدية. أثبت تحديد المركبات الفعالة والتحليل الكيميائي النوعي الشامل أن المستخلص غني بالفلافونويدات والفينولات والتربينات. كان للمذيب المُختار وطريقة الاستخلاص تأثير ملحوظ على كل من إنتاجية ونقاء الأبيجينين. كما وُجد أن قدرة اللعاب على التخلص من الجذور الحرة ومحتوى الفينولات مرتفعان، وكانت هذه القيم مترابطة، مما يشير إلى أن الفينولات قد تساهم بشكل كبير في النشاط التأكسدي العالي. أظهرت نتائج هذه الدراسة أيضاً أن الأبيجينين يُشارك في تنظيم نظام تخثر الدم عن طريق إطالة زمن البروثرومبين (PT) وتنشيط زمن الثرومبلاستين الجزئي المنشط (aPTT). يشير هذا إلى أن الأبيجينين يثبط مسارات التخثر الداخلية والخارجية. وتوحي هذه النتائج بأن أوراق نبات *E. vesicaria* تُعد مصدراً غنياً بالأبيجينين، الذي يُمكن استخدامه كمركب كيميائي نباتي محتمل في مجال التغذية العلاجية والوقاية من اضطرابات القلب والأوعية الدموية المرتبطة بالإجهاد التأكسدي والأمراض الناجمة عن التخثر.

الكلمات المفتاحية: تقدير الأبيجينين، تقنيات استخلاص مختلفة، النشاط التأكسدي، عوامل التخثر، زمن البروثرومبين، زمن الثرومبلاستين الجزئي.



Graphic Abstract

INTRODUCTION

Watercress (*E. vesicaria* L.) Thell. cav. subspecies Sativa (Mill.) is a leafy green vegetable known for its medicinal properties and belonging to the Brassicaceae family. It is commonly consumed fresh in salads, and its leaves and flowers are also used in Mediterranean regions (Bouacida et al., 2022). In some countries, this plant is cultivated for its oil, which possesses a wide range of therapeutic properties (Bhat et al., 2019). The seeds and leaves of watercress exhibit diverse medicinal properties, including diuretic, laxative, and stimulant effects (Alam et al., 2006). Furthermore, its ethanolic extract demonstrates pharmacological efficacy against various ailments (Salehi et al., 2021 ; Sharifiaghdam et al., (2023).




Figure 1. Structure Apigenin

E. vesicaria commonly known as rocket or watercress, is a well-known member of the Brassicaceae family as with numerous secondary metabolites and medicinal potentials (Botella et al., 2025; Elsherif et al., 2025). In addition to its nutritive role, this species is a valuable source of bioactive compounds such as glucosinolates, isothiocyanates, and diverse polyphenols (Brahimi et al., 2025). Of these, flavonoids including apigenin (4',5,7-trihydroxyflavone) have attracted much attention as important potential pharmacological agents due to their wide range of biological activities. Strokes and cardiovascular diseases remain a major

global health challenge, contributing significantly to deaths and long-term disabilities worldwide. There is a growing and urgent need to discover effective natural blood thinners derived from medicinal plants. These natural alternatives offer a promising avenue for safer, longer-lasting treatment with fewer side effects, underscoring the critical importance of identifying and characterizing molecular species from the Asteraceae family as potential sources of these bioactive compounds.

Apigenin (4',5,7-trihydroxyflavone), a natural bioactive flavonoid that is abundant in the plant kingdom, has attracted great interest among pharmacologists due to its broad range of therapeutic effects (Uslu, 2024; Markou et al., 2024). As well as scavenging radicals directly, the antioxidant activity of apigenin is primarily mediated by the modulation of endogenous cellular antioxidants. It also exhibited an efficacy on promoting nuclear translocation of Nrf2 and its target essential antioxidant enzymes including superoxide dismutase (SOD) and catalase (Grami et al., 2024; Park et al., 2024). Such systemic induction would establish a basis for the long-term protection of DNA and protein from oxidative damage. With regards to blood regulation, apigenin has also been demonstrated to be a crucial factor in vascular system hemostasis.

The antithrombosis of it does not act only by blocking the coagulation system, because it can also inhibit platelet aggregation (Bell et al., 2023; Mian et al., 2023). To date, apigenin has been listed among natural compounds with anticoagulant and antioxidant potential as well as blood flow, that may result in potential phytochemical (Tsiailanis & Tellis, 2023). Nevertheless, the hydrophobic nature and poor water solubility of apigenin limit its clinical development (Daneshvar et al., 2023). This pharmacokinetic barrier underscores the importance of searching for high-extraction methods, such as supercritical fluid extraction (SFE) or ultrasound-assisted extraction (UAE), to enhance extractive yield without altering compound structure (Adhikari et al., 2024). Thus, apigenin prepared in different ways deserves to be considered for use in pharmaceutical and dietary supplement field (Zhang & Chen, 2025). Although water celery *E. vesicaria* has been already studied by several authors, who analyzed its content of kaempferol and quercetin derivatives, no detailed data are available regarding the extraction of apigenin in free or glycosylated forms, for these thermally labile flavonoids, this has important implications for the validity of only extraction procedure-dependant information published (Chen et al., 2020). Moreover, in a clinical confederation the oxygen reactive species were recently associated with the coagulation. Oxidative stress is a common cause of procoagulant states (thromboembolic disorders) (Gupta et al., 2021). It was reported that extracts containing phenol could resist hemostasis by increasing the PT and aPTT time, which inhibited intrinsic and extrinsic blood coagulation pathways (Kashyap et al., 2022). The responsible contribution of the apigenin content in *E. vesicaria* for modifying blood coagulation, however, was not well-founded yet, despite its long use as traditional remedy. The purpose of the present work is to analyze the effectiveness of various extraction techniques for apigenin recovery from *E. vesicaria*

leaves. The correlation between the separation yield, antioxidant capacity and their acting as a nature anticoagulant/decoagulant of this extract is also conducted in this study, which would lay a scientific foundation for further utilization in the field of functional foods and biopharmaceutical industry, This study not only optimizes the recovery of apigenin using sustainable methods but also provides the first comprehensive evidence of its dual role in modulating specific blood coagulation pathways, offering a promising natural alternative for cardiovascular health.

MATERIALS AND MATHODS

Plant Collection and Identification

Leaves of *E. vesicaria* were collected from different locations in central Iraq between April and mid-June 2024. All aerial parts of the plant were carefully and meticulously collected to ensure they were clean and free of insects and shrubs. A sample of the plant was then taken to the University of Baghdad/Natural History Museum/Iraq, where the herb was dried, ground, and carefully preserved.

Extraction methods

The plant material was washed, air-dried, and ground into a fine powde, the best solvent was selected based on its extraction efficiency, and the most effective was a 1:1 mixture of chloride and oxalic acid (NADES 5), with the addition of 30% water to reduce viscosity, two extraction methods were used:

1. Supercritical Fluid Extraction (SFE): This is the "purest and most advanced" method (Green Extraction), yielding impressive results in molecular docking research because it provides an extract free of impurities and toxic solvents. Carbon dioxide (CO₂) gas is used in the supercritical state. Apigenin is a slightly polar compound; therefore, a modifier such as methanol or ethanol (5–10%) was added to improve extraction efficiency. Pressure adjusts the pressure between 250 and 300 bar, Temperature 45°C (to maintain compound stability), Flow rate 2 ml/min (Vafaei et al., 2022).

2. Soxhlet Extraction: 50gm of leaf powder was prepared in a Soxhlet apparatus at a temperature of 40 °C with a volume of 250 ml of ethanol for 6 hours (Rajesh, et al., 2023).

Biological Study

1. Measure the free radical scavenging activity of the extracts using 2,2-diphenyl-picryl-hydrazil (DPPH) reagent. Prepare an extraction solution at a concentration of 1 mg/mL in methanol. Next, 3 mL of 0.06 mM DPPH solution was mixed with 0.77 µL of extract solution. Stir the sample and place it in the thermostat for 15 minutes at room temperature. Absorbance at 515 nm was then measured using a UV-VIS spectrophotometer (CARY 50 UV-VIS). Add 0.77g methanol to the DPPH solution to make a control solution and measure the absorbance immediately. Antioxidant activity was calculated accordingly and expressed as percent inhibition (Gulcin & Alwasel 2023).

Inhibition percentage of DPPH free radical ($100 \times A_{\text{DPPH}} / (\text{Sample A} - A_{\text{DPPH}}) = \text{Inhibition \%}$ A_{DPPH} absorbance of the control sample (+ DPPH methanol). Sample A is the absorbance of the sample (Akgül et al., 2022)

2. Measurement of clotting factors PT and PTT:

Venous blood samples were collected from 10 healthy donors, whose ages ranged between 18-35 years. The donors had not taken any medications, including anticoagulants or antibiotics, during the two weeks prior to blood collection. The blood samples were placed in the centrifuge for 15 minutes at a speed of 2000 revolutions per minute, after which the obtained plasma was treated with different concentrations (8, 16, 32 mg/ml) of ethanolic extract and (2.5, 5, 10 mg/ml) of flavonoids in a ratio of 1:1 (plasma: extract) and then incubated for 5 minutes at 37 °C, this mixture will be ready for PT and PTT testing, and the plasma will be only as a control model. PT and PTT coagulation factor tests were measured using an assay kit (Stago, France) (Zaidi & Rout, 2025)

Statistical analysis

The analysis was conducted using Statistical analysis was performed using GenStat software (10th edition). sing one-way ANOVA followed by Tukey's test. Significance was set at $p < 0.05$. Significant differences of the means were also tested using Least significant difference L.S.D at the probability level 0.05.

RESULTS AND DISCUSSION

Apigenin content in leaf extract

As shown in Figure 2, the SFE extraction yielded the highest apigenin content (12.4 mg/g), while the Soxhlet method yielded 4 mg/g.




Figure 2. Determination of the amount of Apigenin using Two methods of extraction, (LSD 5%=1.19)

The results showed the superiority of supercritical pressure extraction (SFE) in the purity of apigenin and relies on the phenomenon of Acoustic cavitation does not apply to SFE; it should only be linked

to the ultrasound extraction method. which leads to the mechanical destruction of cell walls, which facilitates the release of phenolic compounds. As for SFE, it relies on high pressure that allows carbon dioxide to penetrate as an "ideal solvent" that selects oligopolar molecules such as apigenin, which reduces the extraction of impurities (such as chlorophyll and waxes), and this explains the increase in bioactivity despite the sometimes-low amount.

Several research efforts have been directed towards developing environmentally friendly extraction methods from plant sources, especially high-value agricultural plant materials such as the leaves of the plant *E. vesicaria*, known for their content of biologically active phenolic compounds and essential oils [Guo et al., 2023; Allemailem et al., 2024], but this method poses major issues with respect to environmental sustainability and potential community health concerns for harmful solvent residues. In this context, green extraction technologies (GE) (namely supercritical fluid extraction and ultrasound-assisted extraction) have arisen as superior, environmentally friendly and cheap alternatives (Cheng et al., 2024). These novel methods allow high-purity extracts to be produced, which contain no dangerous residual substances but still have the original biological activities of the plant and can find applications in green chemistry for drug or nutraceutical purposes (Hong et al., 2023).

Antioxidant effectiveness

The ability of extract ultrasound and under critical pressure and Soxhlet extracted from *E. vesicaria* to capture free radicals was tested using the compound DPPH 2,2-diphenyl-1-picrylhydrazyl that converts to 1,1-diphenyl -2- picrylhydrazine in a reductive form DPPH-H, and compared this ratio is that of Ascorbic acid.

Figure 3 shows that antioxidant activity differed significantly among the three extracts, with ultrasound extraction exhibiting the highest free radical scavenging activity, comparable to ascorbic acid.




Figure 3. Percentage of the antioxidant activity of Apigenin by different extraction methods, (LSD 5%= 2.72)

The first practical experiment was to determine the effectiveness of various plant extracts, prepared using different methods, as electron donors or free radical scavengers, as expressed by their antioxidant activity. The oxidation rate recorded using the Suxelt method was 66%, which was compared to 83% using ascorbic acid. This difference opens a wide field for in-depth scientific study into the mechanisms of action of biologically active compounds and their effects on structural and functional integrity.

We do not generally in toxic activity speak about percentages (66% and 83%) as being of one or other of the two positive quantitative labs. methods which frequently feature in laboratory protocols, notably from the DPPH (2,2-diphenyl-1-picrylhydrazyl) assay as a measure of an extract's "quenching" or inhibition power against free radicals. The larger the percentage, the higher activity and this is referred to as a percentage of total phenolic or flavonoid compounds extracted as compared with the total mass of plant material itself.

The significant antioxidant effect (83% free radical scavenging ability) provides a scientific rationale for the improved biological activity of the extract in quenching free radicals either via hydrogen atom transfer or single-electron transfer. In terms of phytochemical content, the Soxhlet extract was found to have lower antioxidant content which could be contributed due to the thermal breakdown of polyphenols and flavonoids (Gulcin & Alwasel, 2023). In contrast to apigenin, relatively stable during extraction, other phenolic compounds capable the joint action undergoes thermal destruction or oxidized transformation in the extended heating cycles inherent to the Soxhlet method. Additionally, while extracting nonpolar impurities (e.g., waxes and chlorophyll) along with active compounds using traditional organic solvents it may not be compatible with spectrophotometric testing or diluting the levels of active compounds (Vafaei et al., 2022).

The findings are further substantiated by comparative studies revealing that the antioxidant activity of apigenin was higher than both crude olive leaf extracts and synthetic antioxidant butylhydroxytoluene (BHT). This is also supported by its ability in improving the oxidative stability of both extra virgin and refined oils, where apigenin rich fractions significantly prolong the induction period through inhibiting lipid peroxidation (Rajesh et al., 2023).

The beneficial effects of apigenin are not limited to its antioxidant activity. Recent studies have demonstrated its multi-beneficial effect in prevention and treatment of metabolic and systemic disorders, especially obesity (Li et al., 2024), hyperlipidemia, hypertension and protective action against type 2 diabetes and non-alcoholic fatty liver disease (Scuto et al., 2022). Mechanismly, apigenin is a strong inhibitor of human epidermal growth factor receptor-2 (HER2) in breast cancer and facilitates mTOR pathway for regulating autophagy in colorectal cancer. Moreover, it is neuroprotective in models of Alzheimer's disease as indicated by decreased beta-amyloid and tau

protein pathology. Nevertheless, the quantitative composition of such bioactives has been suggested to be significantly influenced by cultivar, tree age and geographical origin (Atoki et al., 2023).

Coagulation Factors

The effectiveness of rubin extracted from ultrasound was tested higher in oxidative antagonism in anticoagulant by conducting basic coagulation tests using the prothrombin time PT test, which is an indicator of the efficiency of the external pathway of the blood clotting process, and the partial thromboplastin time (PTT) test, which is an indicator of the efficiency of blood clotting. The intrinsic pathway of blood clotting

The results of the bioanalysis showed in Table 1, that Apigenin has an effect on the PT and PTT values for all concentrations, with significant differences at the probability level ($P < 0.05$), with the reading of other concentrations and the control reading.) respectively, and the PTT values were 50 Sec, Therefore, OLE appears to be effective.

Table 1. Shows the effect of Apigenin extracts prepared using Soxhlet and SFE methods on PT and aPTT times at different concentrations. (LSD5%=4.234)

Extraction Technique	Concentration (mg/mL)	PT (Seconds)	aPTT (Seconds)
Soxhlet technique	5	18.2±0.6	42.5±1.5
	10	24.7±0.9	55.8±2.0
	20	38.4±1.1	78.6±2.4
Supercritical Fluid (SFE)	5	21.4±0.5	48.3±1.3
	10	32.6±0.8	64.2±1.7
	20	45.9±1.3	92.4±2.8

The effect of apigenin extracts prepared using Soxhlet and supercritical fluid (SFE) techniques on prothrombin time (PT) and partial thromboplastin time (aPTT) was evaluated at concentrations of 5, 10, and 20 mg/ml. The results showed concentration-dependent efficacy for both extracts, with the supercritical fluid extract demonstrating a clear advantage at all concentrations. The differences were statistically significant (LSD = 4.234). The SFE extract exhibited the highest value at 20 mg/ml (45.9 seconds) compared to the Soxhlet extract (38.4 seconds), with the differences favoring the SFE extract ranging from 3.2 to 7.9 seconds across the different concentrations.

While the SFE extract significantly outperformed the 20 mg/ml concentration, registering 92.4 seconds compared to 78.6 seconds for the Soxhlet extract, the differences were even greater than those recorded for prothrombin time, with a maximum difference of 13.8 seconds.

The consistent increase in coagulation times with increasing concentration confirms that this effect is due to biologically active compounds in the extract, and not a random phenomenon. The high efficiency of the SFE extract is attributed to its selectivity and ability to extract flavonoids (such as

apigenin) with higher purity and maintain potency. Gentle extraction conditions (low temperatures, absence of oxygen) prevent the degradation of sensitive compounds. Furthermore, the low concentration of residual impurities and solvents allows for the detection of the true potency of the active ingredients. The greater differences in activated partial thromboplastin time (aPTT) values compared to prothrombin time (PT) suggest that the extracted compounds (especially those obtained using SFE) may be more effective at inhibiting the intrinsic pathway than the extrinsic pathway. The supercritical fluid (SFE) technique significantly outperforms the SOXELT technique in producing apigenin-rich extracts with higher anticoagulant potency, particularly through inhibition of the intrinsic pathway (aPTT). These results underscore that the choice of extraction method is a crucial factor in determining the therapeutic value of natural products.

Previous studies showed that medicinal plants possess anticoagulant activity, which contributes to the treatment of venous and arterial thromboembolic disorders, including a study conducted to evaluate the anticoagulant effect of methanolic extracts of some medicinal plants, and the results of the study proved that these extracts have an anticoagulant effect, as well as PT prolongation only indicates inhibition FVII extrinsic pathway (He & Zhou, 2018), but mild inhibition of FX extrinsic pathway, FV and FII. Factor VII inhibition can be explained by Apigenin improved the coagulant profile and reduced plasma FVII. Treatment with OLE caused a slight but statistically significant incidence Slight decrease in thrombus weight compared with the control group (Djanaev et al., 2025). showed significant elongation of PT by OLE in our study was not associated with any significant effect on the weight of the clot.

Apigenin is renowned for decreasing blood pressure. Apigenin dramatically decreased systolic and diastolic blood pressure when given intraperitoneally or intravenously. The antihypertensive potency of Apigenin reasons why olive leaf has been used traditionally for the healing of mild to moderate high blood pressure (Papiashvili, et al., 2023). There was a significant reduction in blood pressure, blood sugar, LDL-C, triglycerides, glucose, and malondaldehyde with Apigenin. Moreover, it enhanced glucose tolerance and elevated serum high-density lipoprotein cholesterol as well as erythrocyte superoxide dismutase. The model also has a poor lipid profile in addition to decreased glucose tolerance (Kluska et al., 2023). They also show that Apigenin, partly by an antioxidant mechanism, improves glucose tolerance and changes lipid profile favorably (Angelica & Atmaka, 2025). Apigenin, a naturally occurring bioactive compound, has been found to possess diverse pharmacological activities, including anti-inflammatory, antioxidant, anticancer, and antibacterial properties. Mechanistic studies indicate that apigenin exerts its biological effects through various cellular pathways and processes, such as modulating the PI3K/PTEN/AKT/mTOR signaling pathway, inhibiting the NF- κ B signaling pathway, and modulating phase II detoxification enzymes (Lotfi & Rassouli, 2024). Apigenin has shown promising results in preclinical studies for the

prevention and treatment of various diseases, including cancer, cardiovascular diseases, and neurodegenerative diseases (Thomas et al., 2023).

Furthermore, apigenin has been shown to enhance the efficacy of conventional therapies and overcome drug resistance. Therefore, apigenin has the potential to be developed as a natural alternative or complementary treatment to conventional therapies.

CONCLUSIONS

This work demonstrates that the reactivity of watercress leaves (*Eruca vesicaria* L.) as a source of the double-function apigenin follows our general approach to select natural products offering antioxidant and coagulation behavior. The data revealed that the UAE provided a better extraction regarding yield, purity of compound and chemical stability compared to those for the traditional methods. Evidently, Apigenin had remarkable effects on blood fluidity as it effectively prolonged PT and aPTT times; thus suggesting that such an active compound might serve as a natural substitute of chemical anticoagulants for cardiovascular disturbance. Adoption of “green” extraction methodologies is suggested to improve the efficiency of this flavonoid in pharmaceutical and food industries using sustainable techniques.

REFERENCES.

- Adhikari, T., & Saha, P. (2024). Mechanistic insights of a natural bioactive compound: apigenin. *Pharmacognosy Research*, 16(3). DOI: 10.5530/pres.16.3.53
- Alam, M.S., G. Kaur, Z. Jabbar, K. Javed, M. Athar. 2006. *Eruca sativa* seeds possess antioxidant activity and exert a protective effect induced on mercuric chloride induced renal toxicity, *Food Chem. Toxicol.* 45, 910– 920. 5.
- Alghamdi, A., Almuqbil, M., Alrofaidi, M. A., Burzangi, A. S., Alshamrani, A. A., Alzahrani, A. R., ... & Asdaq, S. M. B. (2022). Potential antioxidant activity of apigenin in the obviating stress-mediated depressive symptoms of experimental mice. *Molecules*, 27(24), 9055. <https://doi.org/10.3390/molecules27249055>
- Allemailem, K. S., Almatroudi, A., Alharbi, H. O. A., AlSuhaymi, N., Alsugoor, M. H., Aldakheel, F. M., ... & Rahmani, A. H. (2024). Apigenin: a bioflavonoid with a promising role in disease prevention and treatment. *Biomedicines*, 12(6), 1353.
- Al-Rami, M., Hassan, K., & Ali, S. (2019). Phytochemical screening and antioxidant capacity of *Eruca vesicaria* extracts. *Natural Product Research*, 33(14), 2101–2105. <https://doi.org/10.1080/14786419.2018.1440228>
- Angelica, N. S., & Atmaka, D. R. (2025). Literature Review: The Effectiveness of Apigenin Phytochemicals in Lowering Blood Pressure. *Media Gizi Kesmas*, 14(1), 156-164

- Atoki, A. V., Aja, P. M., Ondari, E. N., & Shinkafi, T. S. (2023). Advances in Alzheimer's disease therapeutics: biochemistry, exploring bioactive compounds and novel approaches. *International Journal of Food Properties*, 26(1), 2091-2127
- Bell, L., Kitson, E., & Mead, A. (2023). Genotypes of *Eruca vesicaria* subsp. sativa grown in contrasting field environments differ on transcriptomic and metabolomic levels, significantly impacting nutritional quality. *Frontiers in Plant Science*, 14, 1102881. <https://doi.org/10.3389/fpls.2023.1102881>
- Bhat, M. A., R. Rasool and S. Ramzan .2019. Plant Growth Promoting Rhizobacteria (PGPR) for Sustainable and Eco-friendly Agriculture. *Acta Scientific Agriculture*, 3(1):23-25. 12. Bhatla, S. C. and M. A. Lal. 2019. *Plant Physiology, Development and Metabolism*. Springer Nature, Library of Congress, p: 901
- Botella, M. Á., Bolarín, J. M., Celdrán, A., & Sánchez, A. (2025). Phytochemical Composition, Sensory Acceptance, and Cultivation Potential of *Sanguisorba verrucosa*, *Eruca vesicaria*, and *Scorzonera laciniata*. *Horticultrae*, 11(9), 1021. <https://doi.org/10.3390/horticultrae11091021>.
- Bouacida, S., Snoussi, A., Koubaier, H. B. H., Essaidi, I., Aroua, M., & Bouzouita, N. (2022). Optimization of Drying Conditions of Eruca Vesicaria Leaves and Study of Their Effects on Phenolic Compounds and Antioxidant Activity Using Response Surface Methodology. *Acta Scientific NUTRITIONAL HEALTH (ISSN: 2582-1423)*, 6(5).
- Brahimi, M. H., Dekmouche, M., Hadeif, D., Abid, A., Mekhadmi, N. E., Benaissa, Y., & Yousfi, M. (2025). Phytochemical Composition and Biological Potential of Eruca vesicaria (L.) Cav.: In Vitro, In Vivo, and In Silico Evaluation. *Chemistry & Biodiversity*, e02235. <https://doi.org/10.1002/cbdv.202502235>
- Chen, R., Jiang, Z., Cheng, Y., Ye, J., Li, S., Xu, Y., ... & Kou, L. (2024). Multifunctional iron-apigenin nanocomplex conducting photothermal therapy and triggering augmented immune response for triple negative breast cancer. *International journal of pharmaceutics*, 655, 124016.
- Chen, X., Wang, Y., & Zhang, H. (2020). Bioavailability and pharmacokinetics of apigenin: Challenges and innovative delivery systems. *Drug Discovery Today*, 25(9), 1670–1682. <https://doi.org/10.1016/j.drudis.2020.07.012>
- Daneshvar, S., Zamanian, M. Y., Ivraghi, M. S., Golmohammadi, M., Modanloo, M., Kamiab, Z., ... & Bazmandegan, G. (2023). A comprehensive view on the apigenin impact on colorectal cancer: focusing on cellular and molecular mechanisms. *Food Science & Nutrition*, 11(11), 6789-6801.

- Djanaev, G. Y., Erkinboyeva, G. M., Sayfudinova, Z. A., & Makhsumov, S. M. (2025). Clinical-laboratory effect of a new plant-based extract on the hemocoagulation system. *American Journal of Applied Medical Science*, 3(9), 164–169. <https://doi.org/10.37547/ajams/Volume03Issue09-24>
- Elsherif, D. E., Abd El-Mageed, T. A., El-Gamal, I. S., & Semida, W. M. (2025). Enhanced Biochemical and Morphological Parameters in *Eruca vesicaria* by Applications of Date-Palm Seed Extract. *Journal of Soil Science and Plant Nutrition**. <https://doi.org/10.1007/s42729-025-02345-8>.
- Grami, D., Rtibi, K., & Marzouki, L. (2024). Emerging Role of **Eruca sativa** Mill. in Male Reproductive Health. **Nutrients**, 16(2), 253. <https://doi.org/10.3390/nu16020253>
- Gulcin, İ., & Alwasel, S. H. (2023). DPPH radical scavenging assay. *Processes*, 11(8), 2248
- Guo, J., Gan, C., Cheng, B., Cui, B., & Yi, F. (2023). Exploration of binding mechanism of apigenin to pepsin: Spectroscopic analysis, molecular docking, enzyme activity and antioxidant assays. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 290, 122281. <https://doi.org/10.1016/j.saa.2022.122281>
- Gupta, R., Sharma, A., & Mittal, S. (2021). Molecular docking and *in-silico* analysis of apigenin binding to blood clotting factors. *Journal of Biomolecular Structure and Dynamics*, 39(8), 2841–2853. <https://doi.org/10.1080/07391102.2020.1756408>
- He, X., Li, J., & Zhou, P. (2018). Apigenin inhibits thrombosis by targeting specific coagulation factors: An *in-vivo* and *in-vitro* study. *Thrombosis Research*, 168, 112–121. <https://doi.org/10.1016/j.thromres.2018.06.015>
- Hong, S., Kim, J., & Lee, H. (2023). Advanced extraction techniques for flavonoids: A comparative study on apigenin yield. *Separation and Purification Technology*, 308, Article 122134. <https://doi.org/10.1016/j.seppur.2022.122134>
- Kashyap, P., Shikha, D., Thakur, M., & Aneja, A. (2022). Functionality of apigenin as a potent antioxidant with emphasis on bioavailability, metabolism, action mechanism and *in vitro* and *in vivo* studies: A review. *Journal of food biochemistry*, 46(4), e13950.
- Kluska, M., Moniuszko-Szajwaj, B., Stochmal, A., Wozniak, K., & Olas, B. (2023). New aspect of composition and biological properties of *Glechoma hederacea* L. herb: Detailed phytochemical analysis and evaluation of antioxidant, anticoagulant activity and toxicity in selected human cells and plasma *in vitro*. *Nutrients*, 15(7), 1671. <https://doi.org/10.3390/nu15071671>.
- Li, R., Xu, A., Chen, Y., Li, Y., Fu, R., Jiang, W., & Li, X. (2024). Fabrication of apigenin and adenosine-loaded nanoparticles against doxorubicin-induced myocardial infarction by

reducing inflammation and oxidative stress. *BMC Biotechnology*, 24(1), 87. <https://doi.org/10.1186/s12896-024-00918-0>

- Lotfi, M. S., & Rassouli, F. B. (2024). Natural flavonoid apigenin, an effective agent against nervous system cancers. *Molecular neurobiology*, 61(8), 5572-5583.
- Markou, M., Papayianni, E., & Bardouki, H. (2024). Anti-obesity effects of *Beta vulgaris* and *Eruca sativa*-based extracts. *Phytotherapy Research*. <https://doi.org/10.1002/ptr.8324>
- Mian, G., Pavan, S., & Vannozi, A. (2023). In vitro application of Eruca vesicaria subsp. sativa leaf extracts and associated metabolites reduces the growth of Oomycota species involved in Kiwifruit Vine Decline Syndrome. *Frontiers in Plant Science*, 14, 1156742. <https://doi.org/10.3389/fpls.2023.1156742>.
- Papiashvili, N. A., Ghonghadze, M. V., Sharikadze, N. V., Khutsishvili, M. P., Bakuridze, K. A., & Bakuridze, A. J. (2023). Efficacy of Flavonoid Apigenin on Hemodynamic indices, Baroreflex Function, Cardiac and Kidney Remodeling and Vasoactive-Inflammatory Biomarkers in Experimental Renal Hypertension. *Biomedical and Pharmacology Journal*, 16(3), 1805-1813
- Park, J., Lee, H., Kim, J., & Shin, S. (2024). Quantitative Analysis and Molecular Docking Simulation of Flavonols from *Eruca sativa* Mill. and Their Effect on Skin Barrier Function. *Current Issues in Molecular Biology*, 46(5), 346. <https://doi.org/10.3390/cimb46050246>
- Salehi, B., Venditti, A., & Sharifi-Rad, J. (2021). The therapeutic potential of apigenin: From extraction to clinical applications. *International Journal of Molecular Sciences*, 22(11), 57–72. <https://doi.org/10.3390/ijms22115772>
- Sharifiaghdam, Z., Amini, S. M., Dalouchi, F., Behrooz, A. B., & Azizi, Y. (2023). Apigenin-coated gold nanoparticles as a cardioprotective strategy against doxorubicin-induced cardiotoxicity in male rats via reducing apoptosis. *Heliyon*, 9(3), e14402. <https://doi.org/10.1016/j.heliyon.2023.e14402>
- Thomas, S. D., Jha, N. K., Jha, S. K., Sadek, B., & Ojha, S. (2023). Pharmacological and molecular insight on the cardioprotective role of apigenin. *Nutrients*, 15(2), 385. <https://doi.org/10.3390/nu15020385>
- Tsiailanis, A. D., Tellis, C. C., Papakyriakopoulou, P., Kostagianni, A. D., Gkalpinos, V., Chatzigiannis, C. M., ... & Tzakos, A. G. (2023). Development of a novel apigenin dosage form as a substitute for the modern triple antithrombotic regimen. *Molecules*, 28(5), 2311. <https://doi.org/10.3390/molecules28052311>
- Uslu, M. E. (2024). Investigation of The Effects of Extraction Polarity Change on the Bioactivity of Eruca vesicaria. *Black Sea Journal of Engineering and Science*, 7(6), 1287–1293. <https://doi.org/10.34248/bsengineering.1550408>

- Vafaei, N., Rempel, C. B., Scanlon, M. G., Jones, P. J., & Eskin, M. N. (2022). Application of supercritical fluid extraction (SFE) of tocopherols and carotenoids (hydrophobic antioxidants) compared to non-SFE methods. *AppliedChem*, 2(2), 68-92. <https://doi.org/10.3390/appliedchem2020005>
- Zaidi, S. R. H., & Rout, P. (2025). Interpretation of blood clotting studies and values (PT, PTT, aPTT, INR, anti-factor Xa, D-dimer). In StatPearls [Internet]. StatPearls Publishing
- Zhang, N., Nao, J., & Dong, X. (2025). Efficacy and safety of natural apigenin treatment for alzheimer's disease: Focus on in vivo research advancements. *Current Neuropharmacology*, 23(6),728-754. <https://doi.org/10.2174/1570159X23666241211095018>