

The effect of Aqueous some plants Extract on *Giardia lamblia* in vitro

I. N. Al-kaissi

Collage of Agriculture\ University of Anbar

Abstract

Six plants were purchased from herpelists shopes local. These extracts are the *Punica granetum*; *Artemisia campestris*; *Eucalyptus camaldulensis*; *Cuminum cyminum*; *Mangifera indica*; and *Achillea santolina*.

The serial double fold dilution was done to each extract in sterile labeled tubes after preparation. The stock solutions used were (2000, 1000, 500, 125, 62.5, 31.5) mg/ml. Metronidazole (flagyl) was used as standard drug at concentration of 5ml\125mg. The concentrations tested were (500, 250, 125, 62.5, 31.5, 15.75, 7.87) mg\5ml. Viability was assessed by number of viable parasite; number of dead parasite per-microscopically field. The results showed that *punica granetum*; *Artemisia campestris*; *Eucalyptus camaldulensis*; and *Achillea santolina* revealed significant decreases in number of *Giardia lamblia* trophozoites. While the *Cuminum cyminum* and *Mangifera indica* not appeared remarkable effects in number of *Giardia Lamblia* trophozoites.

تأثير المستخلص المائي لبعض النباتات على الجيارديا لامبليا في المختبر

إبراهيم نواف القيسي

كلية الزراعة/ جامعة الأنبار

الخلاصة

سنة نباتات تم شرائها من محلات العشابين في السوق المحلية. وعملت منها مستخلصات نباتية وهذه المستخلصات هي قشور الرمان *Punica granetum* ونبات الشيح *Artemisia campestris* ، كالبتونز *Eucalyptus camaldulensis* ، الكمون *Cuminum cyminum* ، عنبه *Mangifera indica* ، القيصوم *Achillea santolina*. إستعملت طريقة التخفيف المضاعف والمتسلسل (serial double fold dilution) لكل مستخلص محضر في أنابيب مؤشرة بعد التحضير. المحاليل القياسية هي بتركيز (2000، 1000، 500، 125، 62.5، 31.5) مايكروغرام لكل مليلتر حجم. الدواء القياسي (المسيطر control) المستعمل هو محلول المضاد الحيوي metronidazole (flagyl) بتركيز 5\125 mg. التراكيز المختبرة للمسيطر (control) هي (500، 250، 125، 62.5، 31.5، 15.75، 7.87) ملي غرام لكل 5 مليلتر حجم. الحيوية تقاس بواسطة تعداد الطفيليات الحية وكذلك تعداد الطفيليات الميتة في الحقل المجهرى. وظهرت النتائج ان قشور الرمان (الدباغ) *punica granetum*، ونبات الشيح *Artemisia campestris* وقلق الكالبيتونز *Eucalyptus camaldulensis* و*Achillea santolina* أوضحت نقص معنوي في عدد المتغذيات اللامبلية *Giardia lamblia* trophozoites بينما نبات الكمون *Cuminum cyminum* ونبات العنبه *Mangifera indica*. لم تظهر تأثيرات معنوية واضحة في عدد المتغذيات اللامبلية *Giardia lamblia trophozoites*.

Introduction

Giardiasis is a protozoan infection principally of the upper small intestine; it can remain asymptomatic; bring on acute self-limited diarrhea. It lead to intestinal symptoms such as chronic diarrhea, steatorrhea, abdominal cramps, bloating, frequent loose and pale greasy stools, fategne, malabsorption of fats and weight loss.

Infection agent is *Giardia lamblia* a flagellate protozoan. Its occurrence a world wide. Children are infected more frequently than adults. Prevalence is higher in area of poor sanitation and in institutions with children not toilet trained. The prevalence of stool positivity in different areas may range between 1% and 30% depending on community and age group surveyed.

The reservoir is humans; possibly beaver and other wild and domestic animals. The mode of transmission is person to person occurs by hand to mouth transfer of cysts from feces of an infected individual. Concentrations of chlorine used in routine water treatment do not kill *Giardia* cycts, especially when the water is cold. Unfiltered stream and lake waters open to contamination by human and animal feces are a source of infection (1).

Jokipii and Jokipii (2) indicated the need for completely new class of drug for *Giardia* treatment due to the suspected carcinogenicity of metronidazole. Julein, (3); Wolfe, (4), considerable variation in the sensitivity of different stocks of *Giardia lamblia* to the common drug metronidazole furazolidone (5), (6) and treatment failure (7).

Singh et,al (8) reaveled that in modern medicine also plants occupy avery significant place as a raw meterial for some important drugs, although synthetic drugs and antibiotics brought about a revolution in controlling different disease. But these synthetic drugs are out of reach of million of people. Those who live in remote places depend on traditional healers, whom they know and trust, judicious use of medicinal herbs can even cure deadly disease that have long defied synthetic drugs.

The main aim of the present study is to evaluate the *in vitro* anti giardial activity of six crude extract of different Iraqi medicinal plant that were used previously in folk medicine in the treatment of giardiasis and other parasitic disease to examine their traditional uses and their potential to be used in the future as a source for the discovery of new drugs for the treatment of giardiasis in human and animals.

WHO (9) recommended that research shoulel continue for safe, cheap, early administered drug, bearing in mind the need also for effective treatment of patient who are unresponsive to current first line drugs. It was reported that, medicinal plant either in the form of crude drugs or their isolated medicinally active substances, will be important among the treatment choices available to physicians (10).

As conclusion, non of drugs described above can be used with completed assured safety especially to the fetus (11). There are wide varieties to medicinal plants which have pharmacological actuality against several human now (12).

Among them, many plant with antiparasitic spectrum like, *Eucalyptus comaldulensis*, *punica granctum*, *mangifera indica*, *caminum cynminum*, *Artemisia campestris* and *Achillea santolina*, which have good activity against diarrhea and dysentery (13, 14). It was reported that the plant, which contain berberine compounds, may be useful against *Giardia lamblia* (15). Because of less side effect of the plants extracts in comparison with chemical drugs. The investigatores approach to this field now days (16).

Materials and Methods

- Parasite:

Samples collection from individuals complaining of diarrhea were examined for *Giardia lamblia* infection. These patients attend General hospital in Ramadi; consulting laboratory at the college of Medicine and private laboratories in Ramadi city during the period of this work in 2008. Samples examination for *Giardia lamblia* trophozoites and cysts were detected by direct fecal wet preparation method according to the method mentioned by Henandez, *et al* (17).

Cysts purification from Cyst-bearing feces were obtained from infected parasites. Cysts purification was done in present work according to Buchel *et al.*, (18). Fecal samples were carefully diluted in distilled water (1:10). Filtered through 4 layers of gauze, then through a 125 mm, 90 mm aperture metallic mesh and through a 20 mm aperture nylon mesh. Four to five ml of filtrate was placed over 3 ml of chilled 0.85 M sucrose solution in a 10ml conical centrifuge tube. After centrifugation at 4000 r.p.m for 5 min. at room temperature, the water –sucrose interface was removed. Diluted (1:10) with distilled water and centrifuged for 5 min. The pellet was resuspended in 4 ml of water and sucrose gradient was repeated. The washing step was repeated at least for three times. Purified cysts suspension were used freshly or stored at 4 c in distilled water until it was needed.

Excystation of *Giardia lamblia in vitro* was done in the present research according to the method of Bingham and Meyer (19); and the method described by Sauch (20). One volume of purified cysts suspension was added to 10 volume of HCL- saline (0.7 ml concentrated HCL 100ml of 0.85M NaCl, PH 1.5). Three of reducing solution (Hanks Balanced salt solution Ph 7.2 was supplemented with 0.098 gm glutathione and 0.1 gm of L-cysteine - Hcl) and 3 ml of solution bicarbonate prepared freshly. The suspension was mixed and incubated at 37 c for 1 hr. The suspension was centrifuged at 4000 rpm for 5 min at room temperature. The pellet was finally suspended in 5 ml of pre-warmed Excystation medium, which was prepared freshly by (0.015 gm NaHco₃ and 0.01 gm of L-cysteine Hcl monohydrate were dissolved in 4 ml of 5% peptone and 1 ml of (HBSS). The final volume was adjusted to 10 ml with distilled water pH7.1 water bath with 0 c. The peptone was prepared by adding 5% weight \volume peptone to distilled water, boiled in stirring for 10 min. and then stored at 4c⁰.

The sample was incubated in the Excystation medium for 30 min at 37c .Samples were removed from the Excystation tube and placed on depression slides and then covered with glass cover slip sealed with Vaseline – Position for 30 min at 37c⁰.

Cultivation in the present research, attempts for cultivating *Giardia lamblia* trophozoites excysted *in vitro* by using three types of modified media (karapetyan; TYIS-33 and HSP-1) through a several attempts according to the study mentioned by Al-Kaissi (21).

- Plant materials:

Six plants were purchased from herpelist shops in local market. Selection of plants was based on ethobotanical information either literature or directly by personal communications with famous herpelist in Ramadi city. The identification of selective plants was kindly confirmed by Dr. Tarek. M.AL-Fahdawi (profesore of Medicinal plants) Collage of Agriculture\ University of Anbar. These studies was done in the laboratories at college of Medicine, University of Anbar. These plants were: Cuminum cyminum (Umbelliferae), an annual herb with thread like leaves, small white or rose-coloured flowers in compound umbels. Seeds are similar to caraway (22). The plant contains protein; Ca; P; Fe; carotene; Vit. A; Vit. C; fixed oil; volatile oil rich in caminic aldehyde; alpha and B-pinene; Dipantene; phellondrene; Part used: Fruits

known as seeds (13). *Punica granatum*: (punicaceae), a large tree 3-4 m high. Leaves are shiny green. Flowers are large bright orange to red. Fruit are globose to sub globose 8-12 cm in diameter with persistent tubular calyx and leathery skin. The rind is very rich in tannin and contains the alkaloid pelletierine C₈H₁₅ON (22). The fruit rind is good for diarrhoea and dysentery. Part used fruit rinds (13). *Mangifera indica*: (Anacardiaceae). The fruit contains cellulose; yellow coloring matter; Vit. C ; sugar; CH₃; albuminoids; *Mangiferin*, the bark contains tannin and the kernel. Contains tannin; gallic acid; sugar ; fat; gum; starch. Part Used are fruits (13). *Artemisia campestris* (campositae): A wild herb 30-50cm high with many branching stems and ovate or biculate leaves, upper leaves are smaller and in clusters. The plant contains thujone, santonin, the sterols betasitosterol and stigmasterol (14). It has an anti-hemorrhagic activity. Part Used are whole plant.(13).

- ***Eucalyptus camaldulensis* (myrtaceae):** A high tree with grey deciduous bark and narrow lanceolate leaves. flowers 4-8 together are in stalked umbel. Fruit capsule is nearly glabrous. The plant contains mostly kinotannic acid, kinain, kinored, catechin and pyrocatechin (14). It used as astringent in diarrhea and dysentery. Part used are Leaves, flowers and bark (13).
- ***Achillea santolina* (compositae):** It is a small perennial herb. The ray florets are yellow and very short and the herb is hairy having a fragrant odour. the plant contains volatile oil containing azulenes, terpenes, sesquiterpene lactones, choline and glycine (14). herb possesses insect repellent properties. Leaves against dysentery; intestinal colic; expulsion of gases. Part used are The entire herb (13).
- **Plant Extraction:** Each of six plants were dried at room temperature garbled and powdered by rotel coffee grinder type 24 (22). 20 grams of the resulting powder from each plant were mixed with (500)ml distilled water by using electrical blender for (10) min. The suspension was kept shaking over night at room temperature, then centrifuged at 4000 rpm for 5 min and the centrifuges was dried under vacuums by rotary evaporator. The dried extracts were kept in deep freeze (-20 c) until the time of used (23).
- **Astock solution:** The stock solution of each plant was done by dissolving 200 mg of the extract in 100 ml of (HBSS) to get a concentration of 2000 mg/ml which was the highest concentration tested. Sterilization was done by filtration wares through a Millipore 0.45 mm and 0.22 mm. Serial double fold dilution was done to each extract in sterile labelled tubes. The concentration tested were (2000, 1000, 500, 125, 62.5, 31.25) mg/ml (24,25).
- **Standard drug:** Metronidazole (flagyl) was used as standard drug at a concentration of 5 ml 125mg the concentration tested were (500, 250, 125, 62.5, 31.25, 15.75, 7.87) mg/ml Viability was assessed as number of viable parasite, number of dead parasite per microscopic field. The percentage of viability was calculated as below:-

$$\% \text{Viability} = \frac{\text{No. living parasi/ml}}{\text{Total No. ml}} \times 100$$

(The mean of ten trials was used)

Results and Discussion

- **Cultivation:** Using the *In vitro* Excystation procedure sample of cysts from patients with symptomatic and asymptomatic giardiasis were inoculated on manufactured media in the laboratory. These media were (karapetyan, 1962 medium; TYIS-33 medium, keister, 1983; and HSP-1 medium, Meyer, 1976) (21). Although the parasite failed to multiply for (72-hrs) in karapetyan, (1962) they live throughout the whole period of work. The parasites were maintained for (96 hrs.) as TYIS-33 media.
- **Effect of crude plants on *Giardia lamblia* trophozoites:** Some of the crude extracts of six Iraqi plants showed anti-giardial activity against *Giardia lamblia* trophozoites *in vitro*. Our results showed that *punica granatum*; *Artemisia campestris*; *Eucalyptus comaldulensis*; and *Achillea santolina* revealed significant decreases in number of *Giardia lamblia* trophozoites, while the *Cuminum cyminum* and *Mangifera indica* not appeared remarkable effects in number of *Giardia lamblia* trophozoites. It appeared figures below:

Table (1) shows viability of *Giardia lamblia* parasites in *cuminum cyminum* plant at percentage 25% viability with concentration 2000 mg/ml. This plant showed less effect on the density of parasite trophozoites. Table (2) shows viability of *Giardia lamblia* parasites in *Punica granatum* plant. This plant showed a high effect on the density of parasite trophozoites. The result showed the best activity of this plant. Table (3) shown that the aqueous extract of *mangifera indica* had lowered statistically ($P < 0.05$) effect even at 2000mg/ml. Table (4) shows viability of *Giardia Lamblia* parasites in *Artemisia campestris* plant. The aqueous extract of this plant had good effect on trophozoites parasite. Table (5) revealed that viability of *Giardia lamblia* trophozoites were remained unchanged at concentration (31.25), (62.5) mg/ml. although the effect of *Eucalyptus camaldulensis* plant an *Giardia lamblia* trophozoites were finally good effectation. Table (6) shows viability of *Giardia lamblia* parasites in *Achillea santolina* plants. That plant showed good effect on *Giardia lamblia* trophozoites, and the effectation began from 62.5con.mg /ml (viability 87%). The viability was 0% at concentration 2000 mg/ml. The results of mine studies had been showed the effect of aqueous plants extract on *Giardia lamblia* trophozoites *in vitro*. Up to mine knowledge, there were no studies in Iraq concerning anti-giardial activity of medicinal plants. So that, this study in from for the first time that Iraqi medicinal plants have an effect against *Giardia lamblia* parasites although these parasites are fastidious .The lowest concentration (31.25 mg/ml) showed no activity anal forms in a comparison with controls. The on way analysis of variance of the results on six plants indicated that two well separated groups of plants were found. The first group represents *Caminum cyminum* and *Mangifera indica*. These plants demonstrated no significant activity ($P > 0.05$) against the parasite. This result agrees with Al-Wahach (26). The highest concentrations (2000 mg/ml) were eliminated all the parasites and showed good activity in all forms represented the second group consist of *Punica granatum*; *Artemisia campestris*; *Eucalyptus comaldulensis*; and *Achillea santolina*. These plants revealed significant activity ($P < 0.05$) against *Giardia lamblia* trophozoites. These viability percentage at concentration (1000mg/ml) showed (25%); (28.5%); (42.8%); and (57%). The highest concentrations eliminated all the parasites, where no considerable activity was observed at the lowest concentrations. *Punica granatum* had good activity for diarrhea and dysentery (13). It contains the Alkaloids pelletierine (14). *Artemisia campestris* contains santonin, resin, abinthin and thujone and also it was reported to yield camphor (27). Hajawi , *et al* (28) reported that plant *Eucalyptus comaldulensis* contains ecalyptol; phellanderene and pinene. Al-sawah (29)

was reported that the plant *Achillea santolina* contains sesquiterpene lactones which has an insecticidal activity.

The usage of plants may be with less toxicity as reported by Al-Jeboory (30) (drugs isolated from plants are generally less toxic than synthetic drugs), while all available literature agrees that all drugs currently used in the treatment of *Giardia lamblia* are carcinogenic and expensive (31), on the other hand, the procedure of extraction that were adopted in the present study is not too costly since we needed distilled water only as solvents in addition to the non-costly types of plants and their availability in local market. Plants had played an important role in the history of humanity for providing crude drugs along the past centuries. In addition the natural plants drugs served as useful prototypes for even better medicine (32). According to the above results which recorded an *in vitro* anti-giardial activity in some extracts of Iraqi medicinal plants, and the *in vitro* activity may be used as a rough predictor *in vivo* activity (33). Thus, it is concluded that these plants are promising and important for the future treatment of giardiasis and research should continue to prove their *in vivo* activity and their safety for humane use without side effects in order to achieve the aim of using the plants extracts safely and effectively in the treatment of Giardiasis and research should continue to prove their *in vivo* activity and their safety for human use without side effects. In order to achieve the aim of using the plants extracts safely and effectively in the treatment of human and animals giardiasis I am like to recommend that the use of laboratory animals, infected with *Giardia lamblia*.

And studying the effect of the crude extracts in the treatment of these infected animals. Further studies, including sub-fractionation of the extracts using appropriate pharmacognostic method such as chromatography to isolate, identify, and study the active compounds as aphytotherapy, to facilitate their used in human and animals (34).

The data obtained from the present study demstrated *in vitro* that the a queaus extract of some plants on viability percnrntage of *G-lamblia* trophozaites in differentes concentrations

| Plants | 31.25 mg/ml %viable | 62.5 mg/ml %viable | 125 mg/ml %viable | 250 mg/ml %viable | 500mg/ml %viable | 1000mg/ml %viable | 2000mg/ml %viable | Conc. %viable |
|-------------------------------------|------------------------|-----------------------|----------------------|----------------------|---------------------|----------------------|----------------------|------------------|
| Cominum Cyminum | 100% | 100% | 100% | 85.7% | 85.7% | 62.5% | 25% | |
| Punica Granetum | 100% | 87.5% | 75% | 62.5% | 50% | 25% | 0 | |
| Magifera Indica | 100% | 100% | 100% | 100% | 100% | 87.5% | 75% | |
| Artemisia Campestris | 100% | 87.5% | 75% | 57% | 50% | 28.5% | 0 | |
| Eucalyptus Camaldulensis | 100% | 100% | 85.7% | 71.4% | 57% | 42.8% | 0 | |
| Achillea Santolina | 100% | 87.5% | 75% | 71.4% | 66.6% | 57% | 0 | |

Table (1) The mean viability of *Giardia Lamblia* in *Caminum Cycinum*

| 31-25 mg/ml | | | | 62 mg/ml | | | | 125mg/ml | | | | 250mg/ml | | | |
|-------------|---|---|------|-----------|---|---|------|-----------|---|---|-----|---|---|---|------|
| V | d | T | % | V | d | T | % | V | d | T | % | V | d | T | % |
| 8 | 0 | 8 | 100 | 8 | 0 | 8 | 100 | 8 | 0 | 8 | 100 | 6 | 1 | 7 | 85.7 |
| 500mg/ml | | | | 1000mg/ml | | | | 2000mg/ml | | | | v=viable d=dead T=total %=percentage | | | |
| V | d | T | % | V | d | T | % | V | d | T | % | | | | |
| 6 | 2 | 8 | 85.7 | 5 | 3 | 8 | 62.5 | 2 | 6 | 8 | 25 | | | | |

Table (2) The mean viability of *Giardia Lamblia* in *Punica Grantium*

| 31-25 mg/ml | | | | 62 mg/ml | | | | 125mg/ml | | | | 250mg/ml | | | |
|-------------|---|---|-----|-----------|---|---|------|-----------|---|---|----|---|---|---|------|
| V | d | T | % | V | d | T | % | V | d | T | % | V | d | T | % |
| 7 | 0 | 7 | 100 | 7 | 1 | 8 | 87.5 | 6 | 2 | 8 | 75 | 5 | 3 | 8 | 62.5 |
| 500mg/ml | | | | 1000mg/ml | | | | 2000mg/ml | | | | v=viable d=dead T=total %=percentage | | | |
| V | d | T | % | V | d | T | % | V | d | T | % | | | | |
| 4 | 4 | 8 | 50 | 2 | 6 | 8 | 25 | 0 | 8 | 8 | 0 | | | | |

Table (3) The mean viability of *Giardia Lamblia* in *Mangifera indica*

| 31-25 mg/ml | | | | 62 mg/ml | | | | 125mg/ml | | | | 250mg/ml | | | |
|-------------|---|---|-----|----------|---|---|------|-----------|---|---|-----|---|---|---|-----|
| V | d | T | % | V | d | T | % | V | d | T | % | V | d | T | % |
| 6 | 0 | 6 | 100 | 6 | 0 | 6 | 100 | 7 | 0 | 7 | 100 | 8 | 0 | 8 | 100 |
| 500mg/ml | | | | 1000mg/m | | | | 2000mg/ml | | | | v=viable d=dead T=total %=percentage | | | |
| V | d | T | % | V | d | T | % | V | d | T | % | | | | |
| 8 | 0 | 8 | 100 | 7 | 1 | 8 | 87.5 | 6 | 2 | 8 | 75 | | | | |

Table (4) The mean viability of *Giardia Lamblia* in *Artemisia campestris*

| 31-25 mg/ml | | | | 62 mg/ml | | | | 125mg/ml | | | | 250mg/ml | | | |
|-------------|---|---|-----|----------|---|---|------|-----------|---|---|----|---|---|---|----|
| V | d | T | % | V | d | T | % | V | d | T | % | V | d | T | % |
| 8 | 0 | 8 | 100 | 7 | 1 | 8 | 87.5 | 6 | 2 | 8 | 75 | 4 | 3 | 7 | 57 |
| 500mg/ml | | | | 1000mg/m | | | | 2000mg/ml | | | | v=viable d=dead T=total %=percentage | | | |
| V | d | T | % | V | d | T | % | V | d | T | % | | | | |
| 3 | 3 | 6 | 50 | 2 | 5 | 7 | 28.5 | 0 | 8 | 8 | 0 | | | | |

Table (5) The mean viability of *Giardia Lamblia* in *Eucalyptus camaldulensis*

| 31-25 mg/ml | | | | 62 mg/ml | | | | 125mg/ml | | | | 250mg/ml | | | |
|-------------|---|---|-----|----------|---|---|------|-----------|---|---|------|---|---|---|------|
| V | d | T | % | V | d | T | % | V | d | T | % | V | d | T | % |
| 7 | 0 | 7 | 100 | 8 | 0 | 8 | 100 | 6 | 1 | 7 | 85.7 | 5 | 2 | 7 | 71.4 |
| 500mg/ml | | | | 1000mg/m | | | | 2000mg/ml | | | | v=viable d=dead T=total %=percentage | | | |
| V | d | T | % | V | d | T | % | V | d | T | % | | | | |
| 4 | 3 | 7 | 57 | 3 | 4 | 7 | 42.8 | 0 | 7 | 7 | 0 | | | | |

Table (6) The mean viability of *Giardia Lamblia* in *Achillea santolina*

| 31-25 mg/ml | | | | 62 mg/ml | | | | 125mg/ml | | | | 250mg/ml | | | |
|-------------|---|---|------|----------|---|---|------|-----------|---|---|----|---|---|---|------|
| V | d | T | % | V | d | | % | V | d | | % | V | d | | % |
| 8 | 0 | 8 | 100 | 7 | 1 | 8 | 87.5 | 6 | 2 | 8 | 75 | 5 | 2 | 7 | 71.4 |
| 500mg/ml | | | | 1000mg/m | | | | 2000mg/ml | | | | v=viable d=dead T=total %=percentage | | | |
| V | d | T | % | V | d | T | % | V | d | T | % | | | | |
| 4 | 2 | 6 | 66.6 | 4 | 3 | 7 | 57 | 0 | 7 | 7 | 0 | | | | |

References

1. David, L. H. (2004). Control of communicable Diseases Manual. Copyright by American public Health Association. P. 229-231.
2. Jokipii, A. M. M. & Jokipii, L. (1980). *In vitro* susceptibility of *Giardia lamblia* trophozoites to metronidazole and tindazole. *J. Infect Di.*,141(3):317-325.
3. Julein, R. M. (1988). Drugs and the body. W. H. Freeman and company Oxford. England. P.211.
4. Wolf, M. S. (1992). Giardiasis. *Clin. Microbiol.*, 5(1):93-100.
5. Boreham, P. F. L.; Phillips, R. E. & Shepherd, R. W. (1984). The sensitivity of *Giardia intesnalis* to drug *in vitro*. *J. Antimicrob. Chemother.*, 14:449-461.
6. McIntyre, P. L.; Boreham, P. F. L.; Phillips, R. E. & Shepherd, R. W. (1986). Chemotherapy in giardiasis: clinical response and *in vitro* drug sensitivity of human isolates in axenic culture. *J. Pediatr.*, 108:1005-1010.
7. Boreham, P. F. L.; Phillipe, R. E. & Shepherd, R. W. (1987). Heterogeneity in the responses of clones of *Giardia intestinalis* to anti giardial drugs. *Trans. R. Soc. Trop. Med. Hyg.*, 81:406-407.
8. Singh, M. P.; Oraon, B. C. & Narendra, P. (2009). Medicinal plants. APH publishing corporation. New Delhi, India, P. 350.
9. WHO, (1990). Control of the leishmaniasis. Report of WHO expert. committee- tech. Rep. Series NO.931. Geneva.
10. WHO, (1996). Researching medicinal plants in Turkey. Essential drug monitor, WHO action programme on essential drugs NO.22.
11. Zeibig, E. A. (1997). Clinical parasitology; Apractical approach edited by W.B. Saunders company. USA. P.39-43.
12. Ampofo, O. (1977). The word health organization (WHO), P. 26-30.
13. Majeed, N. K. & Mahmud, M. H. (1988). Iraqi plants and herbs between folk medicine and scientific research (in Arabic) 1st Ed . Baghdad, P. 254.
14. Kotb, F. (1985). Medical plants in Libya .1st ed., Arab. Encyclopedia house, Libya, P.830.
15. Kaneda, Y.; Torii, M.; Tanaka, T. & Aikawa, M. (1991). *In vitro* effects of beriberine sulphate on the growth and structure of *Entamoeba histolytica*, *Giardia lamblia* and *Trichomonas vaginalis*. *Ann. Trop. Med. Parasitol.*, 85:417-425.
16. Shnawa, B. H. (1995). Biological and immunological studies on *Giardia lamblia*. Ph. D. Theses, University of Basrah.
17. Henandez, R. T.; Blasco, C. A. & Sanchez, M. A. (1996). Seasonal prevalences of cryptosporidium and *Giardia* in fections in children attending care centres in salamnca (spain) studied for a period of 15 months. *Eur., J. Epidemiol.*, 12:291-295.
18. Buchel, I. A.; Gorenfolt, A.; Chochillon, C.; Savel, G. & Gobert, J. G. (1987). *In vitro* Excystation of *Giardia* from human. Ascaning electrone microscopy study. *J. Parasitol.*, 73:487- 493.
19. Bingham, A. K. & Meyer, E. A. (1979). *Giardia* Excystation canbe induced in vitro in acidic solution. *Nature (London)*, 277: 301-302.
20. Sauch, J. F. (1988). A new method for Excystation of *Giardia*. *Advance in Giardia research*. P.261-264, ed Wallis, P. M. and Hammoud, B. R. University of college, press. Calgary.
21. Al-Kaissi, I. N. S. (2000). Parasitological and epidemiological study of *Giardia lamblia* infection in Ramadi city during the economic-blockade imposed on Iraq.

22. Tyler, V. E.; Brady, L. R. & Robbers, J. E. (1988). Pharmacognosy. 9th ed., Lea & Fibiger, Philadelphia, P.519.
23. Fournet, A.; Barrios, A. A. & Munoz, V. (1994 a). Leishmanicidal and trypanocidal activity of Bolivian plants. J. Ethnopharmacol., 41:19-37.
24. Bwchel, L. A.; Gorenfolt, A.; Chochillon, C. & Gobert, T.G. (1987). *In vitro* Excystation of Giardia from human. Ascanning electron microscopy study. I. Parasitol., 73:487-493.
25. Bingham, A. K.; Jarroll, E. L. & Meyer, E. A. (1979). Giardia spp. Physical factors of Excystation *in vitro* and Excystation eosin exclusion as determinant of viability. Exp. Parasitol., 47:284-291.
26. Al-Wahach, M. M. (2007). Encyclopedia of plants. Dar Digilla, Aman, Jordan. (in Arabic).
27. Chakravarty, H. L. (1976). Plant wealth of Iraq, a dictionary of economic plants. Ministry of Agriculture, Iraq. 1: 5-43.
28. Hajawi, K.; Al-Masimi, H. H. & Kasim, R. M. J. (2004). Pharmacognosy and Medicinal plants. WWW.daralthaqufa.com, Aman, Jordan. (in Arabic).
29. Al-Sawah, A. L. (1992). The genus Achillea in Iraq. PhD. Thesis, University of Baghdad.
30. Al-Jeboory, A. A. (1994). Nutral pharmacology, the future of medicinal plants in drug and medical in dustry (in Arabia). Dar Al-Hurria press, Baghdad. P.139.
31. Adam, R. D. (1991). The biology of Giardia spp. Reviews: Microbiol., 55(4):706-732.
32. Tyler, V. E.; Brady, L. R. & Robbers, J. E. (1988). Pharmacognosy, 9th ed., Lea & Febiger, Philadelphia, P. 519.
33. Berman, D. J. & Wyler, J. D. (1980). An *in vitro* model for investigation of chemotherapeutic agents in leishmaniasis. J. Infect. Dis., 142(1):83-86.
34. Rudolf, F. W. & Volker, F. (2000). Harbel Medicine. Thieme Stuttgart. New York. P. 438.