

Inhibitory effects of Probiotic on growth and adhesion of some gram negative pathogenic bacteria

Ziad M. Al-Khozai
College of science
Al-Qadisyia University

Abstract:

This work focused on three important human pathogens; *Escherichia coli*, *Salmonella typhi* and *Klebsiella pneumoniae*. Results showed that there is several virulence factors in this bacteria such as capsule, enzymes, motility and fimbriae. *Escherichia coli* included fimbriae type I, II and III. *Salmonella typhi* contained only type II and III. While *Klebsiella pneumoniae* included type I and III. Probiotic preparations from *Lactobacillus acidophilus* appeared to affect the bacterial growth and adhesion. Bacterial growth was inhibited by using stock lactic acid bacteria filtrate and 50:50 diluted filtrate. Higher inhibition zones were recorded during the use of stock filtrate of the probiotic on the pathogenic bacterial isolates. Bacterial adhesion to epithelial cells was inhibited also by using the probiotic. In the case of *Escherichia coli*, the adhesion was reduced from 59-61 to 24-21 and 33-30 bacterium/cell by using the stock and diluted probiotic respectively. In the case of *Salmonella typhi* reduction of bacterial adherence was also observed from 55-53 to 11-13 and 16-14 bacterium/cell by using the stock and diluted probiotic respectively. While in the case of *Klebsiella pneumoniae* from 44-46 to 8-9 and 14-10 bacterium/cell by using the stock and diluted probiotic respectively. This results explained that the bacterial adhesion is a crucial step in the colonization and pathogenesis of bacteria, which can be inhibited by using probiotic preparations.

الخلاصة:

ركز هذا العمل على ثلاث أنواع بكتيرية مهمة ممرضة للإنسان. وهي *Escherichia coli* و *Salmonella typhi* و *Klebsiella pneumoniae*. أوضحت النتائج احتواء هذه البكتيريا على العديد من عوامل الضراوة والاستيطان مثل المحفظة والإنزيمات والحركة والأهداب. أوضحت النتائج احتواء *Escherichia coli* على أهداب من النوع الأول والثاني والثالث، في حين احتوت *Salmonella typhi* على النوعين الثالث والثاني، أما *Klebsiella pneumoniae* فقد احتوت على أهداب من النوع الأول والثالث فقط. كذلك بينت الدراسة بان probiotic المحضر من بكتريا *Lactobacillus acidophilus* قد اثر سلباً على نمو والتصاق البكتريا المرضية. إذ أن النمو البكتيري تم تثبيطه باستخدام راشح النمو البكتيري لهذا الجنس وأيضاً لوحظت نتائج مشابهة عند استخدام الراشح المخفف 50:50، وقد كانت أكثر مناطق التثبيط لوحظت باستخدام راشح بكتريا حامض الحليب LAB غير المخفف على الأجناس المرضية المستخدمة في الدراسة. كذلك تم ملاحظة التثبيط الحاصل على قابلية التصاق البكتريا المرضية على الخلايا الطلائية باستخدام probiotic. ففي حلة بكتريا *Escherichia coli* انخفضت نسبة الخلايا الملتصقة من 59-61 بكتريا/خلية إلى 24-21 و 33-30 بكتريا /خلية عند استخدام المحلول المركز من الراشح (probiotic) والمحلول المخفف منه على التوالي. وفي حالة *Salmonella typhi* كان الانخفاض في أعداد الخلايا الملتصقة من 55-53 بكتريا/خلية إلى 11-13 و 16-14 بكتريا/خلية عند استخدام المحلول المركز من الراشح والمحلول المخفف منه على التوالي. وكذلك بالنسبة *Klebsiella pneumoniae* فقد كان الانخفاض في أعداد الخلايا الملتصقة من 44-46 بكتريا/خلية إلى 8-9 و 14-10 بكتريا/خلية عند استخدام المحلول المركز من الراشح والمحلول المخفف منه على التوالي. هذه النتائج توضح أهمية التصاق البكتريا كخطوة أساسية في عملية الاستعمار والأمراضية والتي يمكن أن تثبط باستخدام مستحضرات probiotic.

Introduction :

The ability of successful pathogens to survive in an immunologically hostile environment is provided by large armamentarium of virulence mechanisms, which includes bacterial evade, neutralize or counter the host defense systems, but also manipulate host homeostasis and normal cell functions(1, 2, 3). Adhesion of bacteria to human tissue surfaces and implanted biomaterial surfaces is an important step in the pathogenesis and infection. Fimbriae (or pili) are a group of

rigid, straight, filamentous appendages on bacterial surface and are often no more than 4 to 7 nm in diameter and from 0.2 up to 20 nm length. Fimbriae composed from protein called pilin, the filamentous nature of fimbriae may mediate the adhesion by adhesins associated with fimbriae (3,4, 5). Probiotics (prebiotics) is a dietary adjuvant that beneficially affects the host physiology by modulating mucosal and systemic immunity, as well as improving nutritional and microbial balance in human intestinal tract. Currently probiotic preparations include different species of LAB (Lactic acid bacteria) mainly (6) . The therapeutic effects of lactic acid bacteria include ; improvement of nutritional quality of food and feed, metabolic stimuli of vitamin synthesis, and enzyme production, stabilization of gut microflora and competitive exclusion of enteric pathogens ,enhancing the innate host defenses by production of antimicrobial substances, reduction of serum cholesterol by assimilation mechanism, decrease risk of colon cancer by detoxification of carcinogens ,and tumor suppression by modulation of cell mediated immunity (7). LAB making large proportion from normal flora of gut and vagina. And demonstrate a wide spectrum of antimicrobial characteristics, including acid and bile resistance, antimicrobial systems(ex: bacteriocin, lactic acid, peroxide) and adhesion to various types of pathogens (8). *Lactobacillus acidophilus* has superior capability of producing lactic acid which is antimicrobial and helps the body protection from harmful bacteria adhering the intestinal mucosa. This bacteria can inhibit the activities of adherence and proliferation of pathogenic bacteria by several ways, such as decreasing luminal pH , rendering specific nutrients unavailable to pathogens, decreasing the redox potential of the luminal environment, and producing hydrogen peroxide under anaerobic conditions and producing inhibitory compounds such as bacteriocin (6, 7). This work was carried in an attempt to investigate the inhibitory rule of probiotic on bacterial growth and adhesion.

Materials and methods

Microbiology and biochemical test:

Three important human pathogens include *Escherichia coli* , *Salmonella typhi* and *Klebsiella pneumoniae* were chosen in this work. Three bacterial isolates for each species ,were obtained from the biology department of college of science at Al-Qadisyia university. These isolates grown on culture media, then biochemical and microbiological aspects were documented, included capsule, hemolysis, lipase, gelatinase, oxidase, motility and catalase according to (10,11, 12).

Fimbriae or Colonization factor antigen (CFA):

Fimbriae type I, II and III were screened according to by determination of agglutination of tanned group A RBCs and bacterial cells in the presence of D-manose (3) .

Probiotic and Bacterial adhesion test

This test included some steps as follow:

Preparation of the probiotic

MRS broth was inoculated with 1% of LAB culture, then incubated anaerobically at 37C° for 24 hours. After incubation the culture was centrifuged at 6000 rpm for 15 minutes, the supernatant was obtained, after adjusting the pH of the filtrate to 6.5 by using NaOH, it was filtered through Millipore filter unit (0.22). This filtrate was used in the next steps (14,15).

Preparation of bacterial suspension:

Ten milliliter of nutrient broth medium was inoculated with bacterial growth, the culture was then incubated at 37C° for over night (O.D.₆₀₀ about 0.4) giving (1*10⁹) cell/ml. Cultures of bacteria were washed twice with PBS (phosphate buffer saline) and centrifuged at 1000g for 20 minutes and resuspended in PBS(14,13).

Probiotic sensitivity assay

This assay was carried by using three bacterial strains for each species and two concentrations of the LAB (lactic acid bacteria) filtrate against the bacterial strains, concentrations are stock filtrate or (concentration-1)and 1:1 dilution with normal saline (concentration-2), result was carried by measuring the minimum inhibition zone by making a 5mm discs by using cork porer (12, 14,15).

Preparation of epithelial cell for probiotic assay :

Uroepithelial cells were collected from the urine of some healthy females by centrifugation at 1000g for 10 minutes the washed three times with PBS and centrifuged at 100g for 10 minutes before resuspending in PBS.

Adhesion Test:

A mixture of 0.2 ml of the bacterial suspension, 0.2 ml of the epithelial cell suspension and 0.1 ml of PBS was incubated at 37 C° for one hour. Unattached bacteria to uroepithelial cells were removed by centrifugation in PBS at 1000g for 10 minutes. The final pellet was resuspended in PBS then a drop of it was put onto a microscope slid, air dried fixed with methanol: acetic acid (3:1) and stained with methylen blue. The adherent bacteria to epithelial cells were observed by compound light microscope. Control of only epithelial cells was included (14).

Results:

Results demonstrate that the bacteria contained some types of fimbriae. Type III fimbriae was found in all bacteria under test, while type II was recorded in *Salmonella typhi* only and type I found in *Escherichia coli* and *Klebsiella pneumoniae* while missing in *Salmonella typhi* as explain in table (1).

Table-1: Illustrate the fimbriae types of the bacteria.

Fimbriae type III	Fimbriae type II	Fimbriae type I	Bacteria
+	-	+	<i>Escherichia coli</i>
+	+	-	<i>Salmonella typhi</i>
+	-	+	<i>Klebsiella pneumoniae</i>

The bacterial inhibition zone were estimated by using three isolates and two concentrations of the LAB probiotic, results showed that both *Klebsiella pneumoniae* and *Salmonella typhi* were more sensitive than *Escherichia coli* especially when treated with stock filtrate or concentration (con. 1)of LAB filtrate ,as explained in table(2).

Table-2: Illustrate the bacterial sensitivity to the probiotic filtrate of the LAB .

Bacteria	zone of inhibition (mm)					
	Isolate number 1		Isolate number 2		Isolate number 3	
	Con. 1	Con. 2	Con.1	Con.2	Con.1	Con.2
<i>Escherichia coli</i>	9	8	11.5	10	12.5	11
<i>Salmonella typhi</i>	13.5	11	14	13.5	13	11.5
<i>Klebsiella pneumoniae</i>	14.5	12	13.5	13	12.5	11.5

Con.1= concentration 1 of LAB (stock bacterial filtrate)

Con.2= concentration 2 of LAB (1:1 of bacterial filtrate: normal saline)

Results of the effect of the probiotic reveled that, the bacterial strains adhesion was reduced by using the same concentrations of the LAB in comparison to control, *Escherichia coli* adhesion to Uroepithelial cells was reduced from 59-61 bacterium/ cell without probiotic to 33-30 and 24-21 bacterium/ cell by using con.2 and 1 of the probiotic respectively as explained in figure (1).




Figure-1 : Illustrate the inhibition of adhesion of *Escherichia coli* on the Uroepithelial cells.

Adhesion of *Salmonella typhi* to Uroepithelial cells was reduced from 55-53 bacterium/ cell without probiotic to 16-14 and 11-13 bacterium/ cell by using con.2 and 1 of the probiotic respectively as explained in figure (2).




Figure-2 : Illustrate the inhibition of adhesion of *Salmonella typhi* on the Uroepithelial cells.

Also the adhesion of *Klebsiella pneumoniae* to Uroepithelial cells was reduced from 44-46 bacterium/ cell without probiotic to 14-10 and 8-9 bacterium/ cell by using con.2 and 1 of the probiotic respectively as explained in figure (3).




Figure-3 : Illustrate the inhibition of adhesion of *Klebsiella pneumoniae* on the Uroepithelial cells.

Discussion:

Although both motile and nonmotile species form biofilms, in motile species, the ability to move using flagella or pili is generally required for efficient cell-to-surface attachment (15,16). The fimbriae and capsule mediate the interactions between the bacterium and its immediate environment. Its importance in promoting the formation of biofilms and stimulate interspecies coaggregation, thereby enhancing the bacterial colonization, and are an essential virulence factors (15,17). Species under study appeared to have several types of fimbriae, these results are similar agreed with the results of previous investigators. Enteropathogenic *E. coli* appear to have predilection of adherence to the host cells is thought to involve bundle-forming pili (4). While *S. typhi* capsular polysaccharide and fimbriae, offer the protection from environmental insults and host non specific immune response facilitating the invasion, by mediating the interaction between the bacteria and the mucus surrounding host epithelial cells (2,16). In *K. pneumoniae*, it has been shown that expression of a polysaccharide capsule is essential for the colonization of the large intestine of mice(4, 15). One interpretation of these data is that capsule and fimbriae is required for initial steps of colonization by interacting with the mucus layers and this step is vital for successful colonization in vivo(3). The concomitant increase in adhesion expression would have the net effect of enhancing bacteria-epithelial cells interactions essential for long-term colonization (17).Inhibitory effects were documented in this work by using the Lactic acid bacteria filtrate , these effects due to secretion of inhibitory substances by this bacteria.Lactic acid bacteria have an inhibitory effect against the gram negative and positive bacteria. Some investigators stated a high inhibitory effects of LAB against the enteropathogenic bacteria (6). While others documented a significant inhibition of LAB on *Proteus mirabilis* , proposing the effect to the presence of active antimicrobial secretions such as lactocidin , plantaricin and acidophilin, these results agreed with the results obtained in this work (9). The obtained data reflect a remarkable decrease in the numbers of all tested adhered bacterial cells to the epithelium cells, these results come in conformity with results obtained by some investigators . This was due to the effect of inhibitory substances found in the filtrate of LAB isolates and the acidic pH which affect the growth of the gram negative bacteria by altering some surface structures (18,19). (20) investigated the effect of *Lactobacillus casei* on *E coli* and found that the inhibitory effect was not caused by bacteriophage of hydrogen peroxide but due to the aggregation of *E. coli* an LAB . (14) reported that precoating of LAB strains reduced the binding of uropathogenic coagulase negative *Staphylococci* and *E. coli* to 8 bacteria / cell . while (9) observed a clear reduction in adhesion of *Proteus mirabilis* after the treatment with LAB filtrates reaching to 3-8 bacteria/ cell. Others found that the biosurfactant surlactin as released by lactobacillus isolates may open the way to the development of anti-adhesive biologic coating against *Enterococcus faecalis* , they reported a decrease in the number of adhering *Enterococcus* reaching to approximately 70% (21,22).

References :

- 1- **Brooks**; Geo F.; Butel; Jane S.; Morse Stephan. A. C.(2001). Medical Microbiology. 26/E. Lange. Medical book. Me Graw: Hill. VSA.
- 2- **Russell**, W.; and Herwald H. (2005). Concepts in bacterial virulence. J. Contributions of Microbiology. Vol 12 .Karger. Switzerland.
- 3- **An**, Y.H. ;and Friedman R. J. (2000). Handbook of bacterial adhesion , principles, methods and applications . Humana press. USA.
- 4- **Groisman**, E.A. (2001). Principles of bacterial pathogenesis. Academic Press. University of California. USA.
- 5- **Anderson**. Roy M.(1993). Medical Microbiology. Mosby. London.
- 6 -**Gibson**, G.R. and Roberfroid, M.R.(2008). Handbook of Prebiotics. CRC. Press .USA.
- 7-**Maitijasic**, B. B.; and Rogelj, I. (1999). Bacteriocinogenic activity of lactobacilli isolated from cheese and baby faeces. J. food technology and biotechnology. Vol. 37(2).
- 8- **Messen**, W.; and De vuyst, L. (2002). Inhibitory substances produced by lactobacilli isolated from sourdoughs – a review. J. Food microbiology. Vol.72(1-2).
- 9- **Al-Jeboury**, G. H. (2005). Probiotic Effect on Adhesion property of *Proteus mirabilis*. M.Sc. thesis in biotechnology. College of science/ Al-Nahrain university.
- 10- **Macfaddin** J.F. (2000). Biochemical tests for the identification of medical bacteria. 3rd ed. The Williams and Wilkins-Baltimor USA.
- 11- **Baron**. Ellen Jo; Peterson Tance. R.; Finegold. Syndey. M.(1994). Diagnostic microbiology 9/E Mosby. USA.
- 12- **Collee**, Gerald.; Fraser Andrew G.; Marmion barrie P.; Simmons. Antony.(1996). Practical Medical Microbiology. 14/E Churchill Livingston. USA.
- 13- **Iwahi**, T. Abc, Y. and Tsuchiya, K. (1982). Virulence of *E. coli* in a sending urinary tract infection in mice. J. Medical Microbiology. 15: 303-316.
- 14- **Wary**, S. K.; S. I H.ull.; S. I. Cook; R. C. Barrish; and Hall, R. A. (1986). Identification and characterization of a Uroepithelial cell adhesion from a uropathogenic isolate *Proteus mirabilis*. J. Inflammatory immunology. Vol. 54 (1).
- 15-**Lewus**, C.B.; Kaiser, A. and Monyville, T. J. (1991). Inhibition of food borne bacterial pathogens by bacteriocins from lactic acid bacteria isolated from meat. J. Applied environmental microbiology., 57:1683-1688.
- 16- **Romeo**, T. (2008). Bacterial biofilms. J. Microbiology and Immunology. Vol.322. Springer USA.
- 17- **Wilson**, M. (2002). Bacterial adhesion to host tissues, mechanisms and consequences . Advances in molecular and cellular microbiology Vol.1. Cambridge university press.UK.
- 18- **Mims**.C.A.; Palfire J.HL.;Roitt I.M.;Derek W.; Rosamurd W.; and Roy, M.A.(1995). Medical Microbiology. Mosby, USA.
- 19- **Chan**, R.C.; Bruce A. W.; and Reid, G. A.. (1984). Adherence of cervical, vaginal and distal urethral normal microflora to human Uroepithelial cells and the inhibition of adherence of gram-negative uropathogenes by competitive exclusion. J. Urology. Vol. 131.
- 20- **Blomberg**, L. H.; A. Conway. (1993). Inhibition of adhesion of *E. coli* k88 to piglet ideal mucous by *Lactobacillus spp*. J. Applied environmental microbiology. Vol.59.
- 21-**Velraeds**, M. M.; Van Der Mei, H. C.; Reid G.; and Busscher, H. J. (1996). Inhibition initial adhesion of uropathogenic *Enterococcus faecalis* by biosurfictants from Lactobacillus isolates. J. Applied environmental microbiology. Vol.62.
- 22-**Lamond**, R.J. (2004).Bacterial Invasion of host cells .J. Advances in molecular and cellular microbiology Vol. 5. Cambridge university press. UK.